

REPORTS IN INFORMATICS

ISSN 0333-3590

Equitable colorings of bounded treewidth
graphs

Hans L. Bodlaender and Fedor V. Fomin

REPORT NO 271

May 2004



Department of Informatics
UNIVERSITY OF BERGEN
Bergen, Norway

This report has URL <http://www.ii.uib.no/publikasjoner/texrap/ps/2004-271.ps>

Reports in Informatics from Department of Informatics, University of Bergen, Norway, is available at
<http://www.ii.uib.no/publikasjoner/texrap/>.

Requests for paper copies of this report can be sent to:

Department of Informatics, University of Bergen, Høyteknologisenteret,
P.O. Box 7800, N-5020 Bergen, Norway

Equitable colorings of bounded treewidth graphs

Hans L. Bodlaender* Fedor V. Fomin†

Abstract

A proper coloring of a graph G is *equitable* if the sizes of any two color classes differ by at most one. A proper coloring is ℓ -bounded, when each color class has size at most ℓ . We consider the problems to determine for a given graph G (and a given integer ℓ) whether G has an equitable (ℓ -bounded) k -coloring. We prove that both problems can be solved in polynomial time on graphs of bounded treewidth, and show that a precolored version remains NP-complete on trees.

1 Introduction

There is a wide believe that almost every natural hard problem can be solved efficiently on graphs of bounded treewidth. Of course this is not true, a nice example is the BANDWIDTH MINIMIZATION problem which is NP-hard even on trees of degree three [11, 25]. Another part of 'folklore' in the Graph Algorithms community is that if some (natural) problem can be solved in polynomial time on trees, one should be able to solve it in polynomial time on graphs of bounded treewidth. However, there are some striking and frustrating examples, like $L(2, 1)$ -COLORING, where an efficient algorithm for trees is known [7], but the complexity of the problem on graphs of treewidth ≥ 2 is still open. For more than ten years, EQUITABLE k -COLORING and ℓ -BOUNDED k -COLORING were also examples of such problems. Both problems can be solved in polynomial time on trees and forests [9, 2, 17], i.e. graphs of treewidth 1 and the existence of a polynomial time algorithm for graphs of treewidth ≥ 2 was an open question. In this paper we introduce the first polynomial time algorithm on graphs of bounded treewidth for both versions of coloring. Due to enormous exponents in the running time, our algorithm is mainly of theoretical interest. Our main technique is quite far from the standard dynamic programming on graphs of bounded treewidth. To convince the reader (and ourselves) that the standard dynamic programming approach is unlikely to be implemented for EQUITABLE k -COLORING on graphs of bounded treewidth, we prove that a precolored version of the problem is NP-complete on trees, hence on graphs of treewidth 1. The main idea behind our polynomial time algorithm is to use recent combinatorial results of Kostochka et al. [21] that allow us to handle graphs with 'large' vertex degrees separately.

Previous results. The EQUITABLE k -COLORING problem has a long history. The celebrated theorem of Hajnal & Szemerédi [13] says that any graph G has an equitable k -coloring for $k \geq \Delta(G) + 1$. This bound is sharp. One of the directions of research in this field was to obtain better upper bounds than $\Delta(G) + 1$ for special graph classes. See the survey [23] for a review of the results in this field.

The k -COLORING problem can be trivially reduced to EQUITABLE k -COLORING problem and thus EQUITABLE k -COLORING is NP-hard. Polynomial time algorithms are known for split graphs [8] and trees [9].

*Institute of Information and Computing Sciences, Utrecht University, 3508 TA Utrecht, The Netherlands, hansb@cs.uu.nl

†Department of Informatics, University of Bergen, N-5020 Bergen, Norway, fomin@ii.uib.no

ℓ -BOUNDED k -COLORING has a number of applications. It is also known as the MUTUAL EXCLUSION SCHEDULING problem (MES) which is the problem of scheduling unit-time tasks non-preemptively on m processors subject to constraints, represented by a graph G , so that tasks represented by adjacent vertices in G must run in disjoint time intervals. This problem arises in load balancing the parallel solution of partial differential equations by domain decomposition. (See [2, 26] for more information.) Also the problems of this form have been studied in the Operations Research literature [3, 22]. Other applications are in scheduling in communication systems [15] and in constructing school timetables [19].

The ℓ -BOUNDED k -COLORING problem can be solved in polynomial time on split graphs, complements of interval graphs [24, 8], forests and in linear time on trees [2, 17]. This is almost all what is known about graph classes where the ℓ -BOUNDED k -COLORING problem is efficiently solvable. When one of the parameters ℓ or k is fixed the situation is different. For example, for fixed ℓ or k the problem is solved on cographs [4, 24] and for fixed ℓ on bipartite graphs [4, 14] and line graphs [1]. For $\ell = 2$ the problem is equivalent to the MAXIMUM MATCHING problem on the complement graphs and is polynomial. Notice that for fixed ℓ the problem can be expressed in the counting monadic second-order logic and for graphs of bounded treewidth linear time algorithm for *fixed* ℓ can be constructed [18]. When ℓ is not fixed (i.e. ℓ is part of the input) even for trees the situation is not simple and the question of existence of a polynomial time algorithm for trees [14] was open for several years.

The problem remains NP-complete on cographs, bipartite and interval graphs [4], on cocomparability graphs and fixed $\ell \geq 3$ [24], on complements of line graphs and fixed $\ell \geq 3$ [10], and on permutation graphs and $\ell \geq 6$ [16]. For $k = 3$ the problem is NP-complete on bipartite graphs [4].

Almost all NP-completeness results for ℓ -BOUNDED k -COLORING for different graph classes mentioned above can also be obtained for EQUITABLE k -COLORING by making use of the following observation.

Proposition 1.1. *A graph G on n vertices is ℓ -bounded k -colored if and only if the graph G' obtained from G by adding an independent set of size $\ell k - n$ is equitable k -colorable.*

Our contribution. A standard dynamic programming approach for the COLORING problem needs to keep $O(w^k n)$ entries, where w is the treewidth of a graph and k is the number of colors. Since the chromatic number of a graph is at most $w + 1$ this implies that the classical coloring problem can be solved in polynomial time on graphs of bounded treewidth. Clearly such a technique does not work for EQUITABLE k -COLORING because the number of colors in an equitable coloring is not bounded by a function of treewidth. For example, a star on n vertices has treewidth 1 and it can not be equitable k -colored for any $k < (n - 1)/2$. One of the indications that the complexity of EQUITABLE k -COLORING for graphs of bounded treewidth can be different from 'classical' is that by Proposition 1.1 and [4], the problem is NP-hard on cographs and thus on graphs of bounded clique-width. (Note that chromatic number is polynomial on graphs of bounded clique-width [20].)

However, one of the properties of equitable colorings making our approach possible is the phenomena observed first by Bollobás & Guy [5] for trees: 'Most' trees can be equitable 3-colored. In other words, for almost all trees the difference between the numbers of colors in an equitable coloring is not 'far' from the chromatic number. Recently Kostochka et al. [21] succeeded to generalize the result of Bollobás & Guy for degenerated graphs and our main contribution — the proof that EQUITABLE k -COLORING can be solved in polynomial time on graphs of bounded treewidth (Section 3) — strongly uses this result. Very roughly, we use the results of Kostochka et al. to establish the threshold when the problem is trivially solved and when it become to be solvable in polynomial time by dynamic programming developed in Section 2. In Section 4 we show that such an approach can not be extended to PRECOLORED EQUITABLE k -COLORING by showing that the precolored version of the problem is NP-hard on trees.

1.1 Definitions

We denote by $G = (V, E)$ a finite, undirected, and simple graph. We usually use n to denote the number of vertices in G . For every nonempty $W \subseteq V$, the subgraph of G induced by W is denoted by $G[W]$. The maximum degree of G is $\Delta(G) := \max_{v \in V} d_G(v)$. A graph G is d -degenerate if each of its nonempty subgraphs has a vertex of degree at most d . A nonempty subset of vertices $I \subseteq V$ is *independent* in G if no two of its elements are adjacent in G .

Definition 1.2. A *tree decomposition* of a graph $G = (V, E)$ is a pair $(\{X_i \mid i \in I\}, T = (I, F))$, with $\{X_i \mid i \in I\}$ a family of subsets of V and T a tree, such that

- $\bigcup_{i \in I} X_i = V$.
- For all $\{v, w\} \in E$, there is an $i \in I$ with $v, w \in X_i$.
- For all $i_0, i_1, i_2 \in I$: if i_1 is on the path from i_0 to i_2 in T , then $X_{i_0} \cap X_{i_2} \subseteq X_{i_1}$.

The *width* of tree decomposition $(\{X_i \mid i \in I\}, T = (I, F))$ is $\max_{i \in I} |X_i| - 1$. The *treewidth* of a graph G is the minimum width of a tree decomposition of G .

Lemma 1.3 (Folklore). *Every graph on n vertices and of treewidth $\leq w$ has a vertex of degree at most w , and has at most wn edges.*

A k -coloring of the vertices of a graph $G = (V, E)$ is a partition A_1, A_2, \dots, A_k of V into independent sets (in which some of the A_j may be empty); the k sets A_j are called the *color classes* of the k -coloring. The *chromatic number* $\chi(G)$ is the minimum value k for which a k -coloring exists. A k -coloring A_1, A_2, \dots, A_k is ℓ -bounded if $|A_i| \leq \ell$, $1 \leq i \leq k$. A k -coloring A_1, A_2, \dots, A_k is *equitable* if for any $i, j \in \{1, 2, \dots, k\}$, $|A_i| - |A_j| \leq 1$.

In the *EQUITABLE k -COLORING* problem, we are given a graph $G = (V, E)$, and an integer k , and we ask whether G has an equitable k -coloring.

Theorem 1.4 ([21]). *Every n -vertex d -degenerate graph G is equitable k -colorable for any $k \geq \max\{62d, 31d \frac{n}{n-\Delta(G)+1}\}$.*

Every graph of treewidth $\leq d$ is d -degenerate and Theorem 1.4 implies the following corollary.

Corollary 1.5. *Every n -vertex graph G of treewidth w is equitable k -colorable for any $k \geq \max\{62w, 31w \frac{n}{n-\Delta(G)+1}\}$.*

2 Covering by equitable independent sets.

Let $S \subseteq V$ be a set of vertices of a graph $G = (V, E)$. We say that S can be *covered by independent sets of sizes $\lceil n/k \rceil$* if there is a set of subsets $A_i \subseteq V$, $i \in \{1, 2, \dots, p\}$, $p \leq |S|$, such that

- (i) For every $i \in \{1, 2, \dots, p\}$, A_i is an independent set;
- (ii) For every $i, j \in \{1, 2, \dots, p\}$, $i \neq j$, $A_i \cap A_j = \emptyset$;
- (iii) For every $i \in \{1, 2, \dots, p\}$, either $|A_i| = \lceil n/k \rceil$, or $|A_i| = \lfloor n/k \rfloor$;
- (iv) $S \subseteq \bigcup_{1 \leq i \leq p} A_i$.

Covering by independent sets is a natural generalization of equitable coloring: A graph G has equitable k -coloring if and only if V can be covered by independent sets of sizes $\lceil n/k \rceil$. We use the following observations in our proof.

Lemma 2.1. *Let $S \subseteq V$ be a vertex subset of a graph G .*

- (a) *If S can not be covered by independent sets of sizes $\lceil n/k \rceil$, the graph G is not equitable k -colorable;*

(b) If S can be covered by p independent sets A_1, \dots, A_p of sizes $\lceil n/k \rceil$ and the graph $G' = G[V - \cup_{1 \leq i \leq p} A_i]$ is equitable $(k-p)$ -colorable, the graph G is equitable k -colorable.

Proof. (a) Let B_1, \dots, B_k be the color classes of an equitable k -coloring of G . Consider the collection of sets $\{B_i \mid 1 \leq i \leq k, B_i \cap S \neq \emptyset\}$, and number them A_1, A_2, \dots .

(b) Use color classes A_1, \dots, A_p , and partition the vertices of G' as in the equitable coloring into color classes A_{p+1}, \dots, A_k . \square

Let G be a graph of treewidth w . The next theorem implies that when the cardinality of $S \subseteq V$ or the number k is at most $f(w)$, where f is a function of w , the question if S can be covered by independent sets of sizes $\lceil n/k \rceil$ can be answered in polynomial time. Because there are graphs that need $\Omega(n)$ colors in an equitable coloring, Theorem 2.2 does not imply directly that for graphs of bounded treewidth the EQUITABLE k -COLORING problem can be solved in polynomial time.

Theorem 2.2. *Let $G = (V, E)$ be an n -vertex graph of treewidth $\leq w$, let S be a subset of V , and let k be an integer. One can either find a covering of S by independent sets of sizes $\lceil n/k \rceil$, or conclude that there is no such a covering in polynomial time when k is bounded by a constant, or when $|S|$ is bounded by a constant.*

Proof. (Sketch.) This can be shown using standard dynamic programming techniques for graphs of bounded treewidth. Note that we can check for a covering of at most $\min\{k, |S|\}$ independent sets of sizes $\lceil n/k \rceil$. An algorithm comparable to those e.g. shown in [6, 27], that also has different table entries / homomorphism classes when sets have different sizes solves the problem in polynomial time on graphs of bounded treewidth. \square

3 Bounded treewidth

The main result of this paper is the following theorem.

Theorem 3.1. *The EQUITABLE k -COLORING problem can be solved in polynomial time on graphs of bounded treewidth.*

Proof. Let $G = (V, E)$ be a graph of treewidth w and let k be an integer. To determine if G has an equitable k -coloring, we consider the following cases.

Case 1. $\Delta(G) \leq n/2 + 1$ and $k \geq 62w$. Since

$$\max_{0 \leq \Delta(G) \leq n/2 + 1} \frac{n}{n - \Delta(G) + 1} = 2,$$

we have that

$$k \geq 62w = \max\{62w, 2 \cdot 31w\} \geq \max\{62w, 31w \frac{n}{n - \Delta(G) + 1}\}$$

and by Corollary 1.5, G is equitable k -colorable.

Case 2. $\Delta(G) \leq n/2 + 1$ and $k \leq 62w$. In this case, it follows from Theorem 2.2 that the question whether G has an equitable k -coloring can be solved in polynomial time.

Case 3. $\Delta(G) > n/2 + 1$. Let $S \subset V$ be the set of vertices in G of degree $\geq n/2 + 2$. By Lemma 1.3, G has at most wn edges, so $|S| \leq 4w$. Thus by Theorem 2.2, it can be checked in polynomial time whether S can be covered by independent sets of sizes $\lceil n/k \rceil$. If S cannot be covered, then by part (a) of Lemma 2.1, G has no equitable k -coloring.

Let $A_i \subset V$, $i \in \{1, 2, \dots, p\}$, $p \leq |S|$, be an equitable covering of S by independent sets of sizes $\lceil n/k \rceil$. We define a new graph $G' = G[V - \cup_{1 \leq i \leq p} A_i]$. The maximum vertex degree in G' is at most $n/2 + 1$ and the treewidth of G' is $\leq w$. Graph G' has

$$n' = |V - \cup_{1 \leq i \leq p} A_i| \geq n - p \left\lceil \frac{n}{k} \right\rceil \geq n - 4w \left(\frac{n}{k} - 1 \right) > \left(1 - \frac{4w}{k}\right)n$$

vertices.

Let $k' = k - p$. We need again case distinction.

Case A. $k' \geq \max\{62w, 31w \frac{n'}{n' - n/2}\}$. Then

$$k' \geq \max\{62w, 31w \frac{n'}{n' - n/2}\} \geq \max\{62w, 31w \frac{n'}{n' - \Delta(G') + 1}\}$$

and by Corollary 1.5, G' is equitable k' -colorable. By part (b) of Lemma 2.1, G has equitable k -coloring.

Case B. $k' < \max\{62w, 31w \frac{n'}{n' - n/2}\}$ and $k' < 62w$. Since $p \leq 4w$, we have that $k = k' + p < 66w$. Then by Theorem 2.2, the question whether G has an equitable k -coloring, can be solved in polynomial time.

Case C. $k' < \max\{62w, 31w \frac{n'}{n' - n/2}\}$ and $k' \geq 62w$. Then

$$k' < 31w \frac{n'}{n' - \frac{n}{2}} < 31w \frac{n}{(1 - \frac{4w}{k})n - \frac{n}{2}} = \frac{31w}{\frac{1}{2} - \frac{4w}{k}}. \quad (1)$$

Using $k = k' + p \geq 62w$, we have that

$$\frac{4w}{k} \leq \frac{4w}{62w} = \frac{2}{31}$$

and

$$\frac{1}{2} - \frac{4w}{k} \geq \frac{27}{62}. \quad (2)$$

By (1) and (2),

$$k' < 31w \frac{62}{27} < 72w$$

and we conclude that $k = k' + p \leq 76w$. Again, by Theorem 2.2 the question if G has an equitable k -coloring, can be solved in polynomial time. \square

By Proposition 1.1, Theorem 3.1 implies directly that there is a polynomial time algorithm for the ℓ -BOUNDED k -COLORING problem restricted to graphs of bounded treewidth.

4 Equitable coloring with precoloring

For a graph $G = (V, E)$, a *precoloring* π of a subset $V' \subset V$ in k colors is a mapping $\pi: V' \rightarrow \{1, 2, \dots, k\}$. We say that a coloring A_1, A_2, \dots, A_k of G *extends* the pre-coloring π if $u \in A_{\pi(u)}$ for every $u \in V'$. We consider the following problem: **EQUITABLE COLORING WITH PRECOLORING:** For a given graph G , integer k and a given precoloring π of G , determine whether there exists an equitable k -coloring of G extending π .

Theorem 4.1. **EQUITABLE COLORING WITH PRECOLORING is NP-complete on trees.**

Proof. We use a reduction from the problem

3-PARTITION

Instance: A set A of non-negative integers a_1, \dots, a_{3m} , and a bound B , such that for all i with $1 \leq i \leq 3m$, $(B + 1)/4 < a_i < B/2$ and $\sum_{1 \leq i \leq 3m} a_i = mB$.

Question: Can A be partitioned into m disjoint sets A_1, A_2, \dots, A_m such that $\sum_{a_i \in A_j} a_i = B$ for every j with $1 \leq j \leq m$?

3-PARTITION is NP-complete in the strong sense (Problem SP15 in Garey & Johnson [12]).

Let the set $A = \{a_1, \dots, a_{3m}\}$ and the bound B be an instance of 3-PARTITION. We construct a tree G and a precoloring of G such that G is equitable $(m+1)$ -colorable if and only if A can be 3-partitioned.

For every $i \in \{1, 2, \dots, m\}$ we define the set $N_i = \{1, 2, \dots, m\} - \{i\}$ and the precolored star S_i as a star with one non-precolored central vertex v adjacent to $m-1$ leaves which are precolored with all colors from N_i . Thus vertex v can be colored only with color i or $m+1$. For every $i \in \{1, 2, \dots, m\}$ and $j \in \{1, 2, \dots, 3m\}$, we define the precolored tree $G_{i,j}$ as a tree obtained by taking the disjoint union of a_j+1 precolored stars S_i and by making the central vertex v of one of them to be adjacent to the central vertices of the others a_j stars. We call the vertex v the central vertex of $G_{i,j}$. Thus $G_{i,j}$ has $m(a_j+1)$ vertices; for every color $\ell \in N_i$ there are (a_j+1) vertices of $G_{i,j}$ precolored with ℓ . Every $(m+1)$ -coloring of $G_{i,j}$ either colors v with $m+1$ and the remaining a_j non-precolored vertices with i , or it colors v with i and the remaining a_j non-precolored vertices with $m+1$. (See Fig. 1.)




Figure 1: Tree $G_{i,j}$. Here $a_j = 3$, $m = 4$, $i = 2$ and $N_2 = \{1, 3, 4\}$. In any 5-coloring of $G_{i,j}$, either v should be colored with 5 and all its non-precolored neighbors with 2, or v should be colored with 2 and the neighbors with 5.

For every $j \in \{1, 2, \dots, 3m\}$, we define a precolored tree G_j as follows: We take the disjoint union of precolored trees $G_{i,j}$, $i \in \{1, 2, \dots, m\}$, add one vertex c_j adjacent to all central vertices of trees $G_{i,j}$ and add one leaf adjacent to c_j precolored with $m+1$. Thus G_j has $m^2(a_j+1) + 2$ vertices; for every color $\ell \in \{1, 2, \dots, m\}$ there are $(m-1)(a_j+1)$ vertices of G_j precolored with ℓ and there is 1 vertex precolored with $m+1$. Also in any coloring of G_j vertex c_j can not be colored with $m+1$. Thus for every $(m+1)$ -coloring of G_j the spectra of colors used on neighbors of c_j does not contain all colors from $\{1, 2, \dots, m\}$.

Finally G is obtained by taking the disjoint union of precolored trees G_1, G_2, \dots, G_{3m} and an independent set of cardinality $3m(m-2) + B$ precolored with color $m+1$, and then making this forest a tree by adding appropriate edges between precolored vertices of the same color in different trees. Thus G has

$$\sum_{1 \leq j \leq 3m} m^2(a_j+1) + 3(m-2) + B = m^2(mB+3m) + 3(m-2) + B$$

vertices. For every color $\ell \in \{1, 2, \dots, m\} - \{i\}$ there are

$$\sum_{1 \leq j \leq 3m} (m-1)(a_j+1) = (m-1)(mB+3m)$$

vertices of G precolored with ℓ and $3m(m-1) + B$ vertex are precolored with $m+1$.

Claim 4.1.1. *A can be partitioned into m disjoint sets A_1, A_2, \dots, A_m such that $\sum_{a_i \in A_j} a_i = B$, if and only if G has an equitable $(m+1)$ -coloring that extends the precoloring.*

Proof. Suppose that A can be partitioned into m disjoint sets A_1, A_2, \dots, A_m such that $\sum_{a_i \in A_j} a_i = B$. We define an extension of the precoloring of G as follows. For every fixed

$j \in \{1, 2, \dots, 3m\}$, we choose i such that $a_j \in A_i$. We color the central vertex of $G_{i,j}$ with color $m + 1$ and the remaining non-colored a_j vertices of $G_{i,j}$ with color i . In each graph $G_{\ell,j}$, $\ell \neq i$, a_j vertices are colored with $m + 1$ and one vertex with ℓ . Also we color vertex c_j with i . Thus in every graph G_j on the set of non-precolored vertices color i is used $a_j + 1$ times. Any color $\ell \in \{1, 2, \dots, m\} - \{i\}$ is used one time and color $m + 1$ is used

$$a_j(m - 1) + 1$$

times on non-precolored vertices. Thus in graph G the number of vertices colored with color $\ell \in \{1, 2, \dots, m\}$ is

$$(m - 1)(mB + 3m) + \sum_{a_j \in A_\ell} (a_j + 1) + \sum_{\{1 \leq j \leq 3m \mid a_j \notin A_\ell\}} 1 = (m - 1)(mB + 3m) + B + 3m.$$

The number of vertices colored with $m + 1$ is

$$3m(m - 2) + B + \sum_{1 \leq j \leq 3m} (a_j(m - 1) + 1) = (m - 1)(mB + 3m) + B + 3m$$

and we conclude that the obtained coloring is an equitable $(m + 1)$ -coloring.

Suppose now that G has equitable $(m + 1)$ -coloring that extends the given precoloring. The main observation here is that for every $j \in \{1, 2, \dots, 3m\}$, at most $a_j(m - 1) + 1$ vertices of a graph G_j are colored with $m + 1$. (Otherwise, the coloring of the central vertices of graphs $G_{i,j}$, $i \in \{1, 2, \dots, m\}$, uses the whole spectra $\{1, 2, \dots, m\}$, thus leaving no space for the color of c_j .) If for some $j \in \{1, 2, \dots, 3m\}$, less than $a_j(m - 1) + 1$ vertices of a graph G_j are colored with color $m + 1$, then (the coloring is equitable) for some $j' \in \{1, 2, \dots, 3m\}$, at least $a_j m$ vertices of the graph $G_{j'}$ are colored with $m + 1$, which is a contradiction.

Thus we can conclude that for every $j \in \{1, 2, \dots, 3m\}$, there is exactly one subgraph $G_{i,j}$ such that a_j non-precolored vertices of $G_{i,j}$ are colored with i . For all other $i' \in \{1, 2, \dots, m\}$, $i \neq i'$, a_j non-precolored vertices of $G_{i',j}$ are colored with $m + 1$. We define

$$A_i = \{a_j \mid a_j \text{ non-precolored vertices of } G_{i,j} \text{ are colored with } i\}.$$

In G , the number of vertices colored with color $i \in \{1, 2, \dots, m\}$ is

$$(m - 1)(mB + 3m) + B + 3m = (m - 1)(mB + 3m) + \sum_{a_j \in A_i} (a_j + 1) + \sum_{\{1 \leq j \leq 3m \mid a_j \notin A_i\}} 1.$$

Thus for every $i \in \{1, 2, \dots, m\}$

$$\sum_{a_j \in A_i} (a_j + 1) = B$$

and A_1, A_2, \dots, A_m is a 3-partition of A . □

So we have a polynomial reduction from 3-PARTITION to EQUITABLE k -COLORING WITH PRECOLORING. As EQUITABLE k -COLORING WITH PRECOLORING trivially belongs to NP, we can conclude it is NP-complete. □

A direct corollary is that ℓ -BOUNDED k -COLORING WITH PRECOLORING is NP-complete for trees.

We can also formulate the related EQUITABLE COLORING WITH PRECOLORING problem. Here, we are given a precolored graph G , and ask whether there exists an integer k , and an equitable k -coloring of G that extends the precoloring. I.e., k is not part of the instance of the problem.

Proposition 4.2. EQUITABLE COLORING WITH PRECOLORING is NP-complete on trees.

Proof. Clearly, the problem is in NP. We transform from EQUITABLE k -COLORING WITH PRECOLORING on trees. Take a tree instance T of EQUITABLE k -COLORING WITH PRECOLORING. We may assume, by the proof above, that for each color i , $1 \leq i \leq k$, T has at least one precolored vertex with color i . Suppose T has n vertices. We construct a tree T' as follows. For each i , $1 \leq i \leq k$, we add $n+1$ new vertices, precolored with i . Each of the new vertices is made adjacent to a precolored vertex in T with a different color. We obtain a new tree T' . T' cannot have an equitable r coloring for some $r \neq k$. (An equitable coloring must use at least k colors. It cannot use more than k colors, as we have at most $n - k$ non-precolored vertices, and at least $n + 2$ vertices are already precolored with color 1.) Also, T' has an equitable k -coloring if and only if T has an equitable k -coloring. (Use the same colors for the non-precolored vertices in both cases.) So, NP-hardness follows. \square

References

- [1] N. ALON, *A note on the decomposition of graphs into isomorphic matchings*, Acta Math. Hungar., 42 (1983), pp. 221–223.
- [2] B. S. BAKER AND E. G. COFFMAN, JR., *Mutual exclusion scheduling*, Theoret. Comput. Sci., 162 (1996), pp. 225–243.
- [3] J. BLAZEWICZ, K. H. ECKER, E. PESCH, G. SCHMIDT, AND J. WEGLARZ, *Scheduling Computer and Manufacturing Processes*, Springer, Berlin, 2001. 2nd ed.
- [4] H. L. BODLAENDER AND K. JANSEN, *Restrictions of graph partition problems. I*, Theoret. Comput. Sci., 148 (1995), pp. 93–109.
- [5] B. BOLLOBÁS AND R. K. GUY, *Equitable and proportional coloring of trees*, J. Combin. Theory Ser. B, 34 (1983), pp. 177–186.
- [6] R. B. BORIE, *Generation of polynomial-time algorithms for some optimization problems on tree-decomposable graphs*, Algorithmica, 14 (1995), pp. 123–137.
- [7] G. J. CHANG AND D. KUO, *The $L(2, 1)$ -labeling problem on graphs*, SIAM J. Discrete Math., 9 (1996), pp. 309–316.
- [8] B.-L. CHEN, M.-T. KO, AND K.-W. LIH, *Equitable and m -bounded coloring of split graphs*, in Combinatorics and computer science (Brest, 1995), vol. 1120 of Lecture Notes in Comput. Sci., Springer, Berlin, 1996, pp. 1–5.
- [9] B.-L. CHEN AND K.-W. LIH, *Equitable coloring of trees*, J. Combin. Theory Ser. B, 61 (1994), pp. 83–87.
- [10] E. COHEN AND M. TARSI, *NP-completeness of graph decomposition problems*, J. Complexity, 7 (1991), pp. 200–212.
- [11] M. R. GAREY, R. L. GRAHAM, D. S. JOHNSON, AND D. E. KNUTH, *Complexity results for bandwidth minimization*, SIAM J. Appl. Math., 34 (1978), pp. 477–495.
- [12] M. R. GAREY AND D. S. JOHNSON, *Computers and Intractability, A guide to the theory of NP-completeness*, W. H. Freeman and Co., San Francisco, Calif., 1979.
- [13] A. HAJNAL AND E. SZEMERÉDI, *Proof of a conjecture of P. Erdős*, in Combinatorial theory and its applications, II (Proc. Colloq., Balatonfüred, 1969), North-Holland, Amsterdam, 1970, pp. 601–623.
- [14] P. HANSEN, A. HERTZ, AND J. KUPLINSKY, *Bounded vertex colorings of graphs*, Discrete Math., 111 (1993), pp. 305–312. Graph theory and combinatorics (Marseille-Luminy, 1990).

- [15] S. IRANI AND V. LEUNG, *Scheduling with conflicts, and applications to traffic signal control*, in Proceedings of the Seventh Annual ACM-SIAM Symposium on Discrete Algorithms (Atlanta, GA, 1996), New York, 1996, ACM, pp. 85–94.
- [16] K. JANSEN, *The mutual exclusion scheduling problem for permutation and comparability graphs*, Information and Computation, 180 (2003), pp. 71–81.
- [17] M. JARVIS AND B. ZHOU, *Bounded vertex coloring of trees*, Discrete Math., 232 (2001), pp. 145–151.
- [18] D. KALLER, A. GUPTA, AND T. SHERMER, *The χ_t -coloring problem*, in Symp. on Theoretical Aspects of Comp. Sc., STACS'95 (Munich, 1995), vol. 900 of Lecture Notes in Comput. Sci., Springer, Berlin, 1995, pp. 409–420.
- [19] F. KITAGAWA AND H. IKEDA, *An existential problem of a weight-controlled subset and its application to school timetable construction*, Proceedings of the First Japan Conference on Graph Theory and Applications (Hakone, 1986), Discrete Math., 72 (1988), pp. 195–211.
- [20] D. KOBLER AND U. ROTICS, *Edge dominating set and colorings on graphs with fixed clique-width*, Discrete Appl. Math., 126 (2003), pp. 197–221.
- [21] A. V. KOSTOCHKA, K. NAKPRASIT, AND S. V. PEMMARAJU, *Coloring d -degenerate graphs equitable*. Manuscript, 2003.
- [22] J. KRARUP AND D. DE WERRA, *Chromatic optimisation: limitations, objectives, uses, references*, European J. Oper. Res., 11 (1982), pp. 1–19.
- [23] K.-W. LIH, *The equitable coloring of graphs*, in Handbook of Combinatorial Optimization, Vol. 3, Kluwer Acad. Publ., Boston, MA, 1998, pp. 543–566.
- [24] Z. LONC, *On complexity of some chain and antichain partition problems*, in Graph-theoretic concepts in computer science WG'91 (Fischbachau, 1991), vol. 570 of Lecture Notes in Comput. Sci., Springer, Berlin, 1992, pp. 97–104.
- [25] B. MONIEN, *The bandwidth minimization problem for caterpillars with hair length 3 is NP-complete*, SIAM J. Algebraic Discrete Methods, 7 (1986), pp. 505–512.
- [26] B. F. SMITH, P. E. BJØRSTAD, AND W. D. GROPP, *Domain Decomposition. Parallel multilevel methods for elliptic partial differential equations*, Cambridge University Press, Cambridge, 1996.
- [27] J. A. TELLE AND A. PROSKUROWSKI, *Algorithms for vertex partitioning problems on partial k -trees*, SIAM J. Discrete Math., 10 (1997), pp. 529 – 550.