

REPORTS
IN
INFORMATICS

ISSN 0333-3590

Finding the Longest Isometric Cycle in a
Graph

Daniel Lokshtanov

REPORT NO 326

June 2006



Department of Informatics
UNIVERSITY OF BERGEN
Bergen, Norway

This report has URL <http://www.ii.uib.no/publikasjoner/texrap/ps/2006-326.ps>
Reports in Informatics from Department of Informatics, University of Bergen, Norway, is
available at <http://www.ii.uib.no/publikasjoner/texrap/>.

Requests for paper copies of this report can be sent to:
Department of Informatics, University of Bergen, Høyteknologisenteret,
P.O. Box 7800, N-5020 Bergen, Norway

Finding the Longest Isometric Cycle in a Graph

Daniel Lokshtanov *

Abstract

When searching for the longest cycle in a graph, we sometimes want to restrict our search space to the cycles that do not have crossing edges - induced cycles. If we think of the crossing edges as shortcuts, looking for induced cycles is then looking for shortcut-free cycles. However, an induced cycle may have a shortcut in the following sense: For two vertices on the cycle, the distance between them in the graph is strictly less than the distance between them in the cycle. As an example, consider the wheel graph on 7 or more vertices. The circumference of the wheel induces a cycle. In this cycle, the distance between diametrically opposite vertices is at least 3. At the same time one can get between any two vertices on this cycle in two steps via the central node. Thus this cycle is induced, but clearly not shortcut-free. We will say that a cycle that has no shortcuts in the above sense is an *isometric* cycle of G . Finding the longest isometric cycle of a graph is then a natural variant of the problem of finding a longest cycle. In this paper we present a polynomial time algorithm for finding the longest isometric cycle in a graph.

1 Definitions and terminology

All input graphs are simple, connected, unweighted and undirected. A *walk* W is a sequence of vertices where each consecutive pair of vertices is connected by an edge. If the first and last vertex of W are the same we say that W is *cyclic*. If all vertices in W are unique we say that the walk is a *path*. If W has at least 3 vertices and all vertices of W are unique, but the first and last vertex are the same, W is a cycle. The *length* of a walk is the number of edges in it. The number of edges in the shortest path between two vertices u and v in a graph G is denoted $d_G(u, v)$ and is called the *distance* between u and v . When the graph is not specified we implicitly mean distance in G and will write $d(u, v)$ for short. If u and v actually are the same vertex, we say that $d_G(u, v) = 0$. If u and v lie in different components of G , the distance between them is infinite. For a natural number p , G to the *power of* p is the graph $G^p = (V(G), \{(u, v) : d_G(u, v) \leq p\})$. A subgraph H of a graph G is an *isometric* subgraph, if for every u and v in $V(H)$, $d_H(u, v) = d_G(u, v)$. Notice that an isometric subgraph is an induced subgraph.

2 A useful auxiliary graph

In this section we are going to concentrate upon an auxiliary graph that we can use to test whether a given graph G has an isometric cycle of length exactly k . We will assume that $k \geq 3$. Using G we build a new graph G_k . The set of vertices of G_k is the set of ordered pairs $\{(u, v) \in V : d(u, v) = \lfloor k/2 \rfloor\}$ and there is an edge between (u, v) and (w, x) in G_k if $(u, w) \in E(G)$ and $(v, x) \in E(G)$. In order to use G_k we

*Department of Informatics, University of Bergen, N-5020 Bergen, Norway. Email: daniel.lokshtanov@student.uib.no.

must say something about the relationship between distances in G and distances in G_k .

Lemma 2.1 $d_{G_k}((u, v), (q, w)) \geq \text{Max}\{d(u, q), d(v, w)\}$.

Proof. Let $P = \{(u, v), (a_1, b_1), (a_2, b_2) \dots (q, w)\}$ be a shortest path from (u, v) to (q, w) in G_k . Then both $P_1 = u, a_1, a_2 \dots q$ and $P_2 = v, b_1, b_2 \dots w$ are paths in G . P , P_1 and P_2 all have the same length, completing the proof. ■

Now, let (u, v) be a vertex in G_k . Notice that such a vertex must exist if there is an isometric cycle of length k in G . The following results will allow us to use G_k as a tool for finding isometric cycles of length k in G .

Lemma 2.2 *If k is even and there is an isometric cycle of length k in G going through u and v then the distance between (u, v) and (v, u) is $k/2$ in G_k .*

Proof. Assume G has an isometric cycle of length k , $C = \{c_1, c_2, \dots, c_{k/2}, c_{k/2+1}, \dots, c_k\}$ with $c_1 = u$ and $c_{k/2+1} = v$. Note that if C contains u and v , but $c_1 \neq u$ we can relabel the vertices of C so that c_1 becomes u . As $d_G(u, v) = k/2$ and C is isometric we know that $c_{k/2+1}$ then must be v . We now know that $(c_1, c_{k/2+1}), (c_2, c_{k/2+2}), \dots, (c_{k/2}, c_k)$ and $(c_{k/2+1}, c_1)$ are vertices in G_k , and there clearly is an edge between each consecutive pair of these vertices. Thus $d_{G_k}[(u, v), (v, u)] \leq k/2$. By Lemma 2.1 we have that $d_{G_k}[(u, v), (v, u)] \geq k/2$ which insures equality and completes the proof. ■

Lemma 2.3 *If k is even and the distance between (u, v) and (v, u) is $k/2$ in G_k then there is an isometric cycle of length k in G going through u and v .*

Proof. Let $d_{G_k}[(u, v), (v, u)] = k/2$ and let $P = \{(u, v), (a_2, b_2), (a_3, b_3) \dots (a_{k/2-1}, b_{k/2-1}), (v, u)\}$ be a shortest path between (u, v) and (v, u) . Now, obviously $W = \{u, a_2, a_3 \dots v, b_2, b_3 \dots, u\}$ is a cyclic walk of length k . By definition of a walk, W is also a subgraph in G . In order to obtain a contradiction let us assume that there is a pair of vertices a and b in W with $d_G(a, b) < d_W(a, b)$. Now, let x be a vertex in W so that either (a, x) or (x, a) is in P . As a and x are on opposite sides of the cyclic walk W , there is a walk of length $k/2$ from a to x going through b . This means that $d_W(a, b) + d_W(b, x) \leq k/2$. But $d_G(a, b) < d_W(a, b)$ implying $d_G(a, x) \leq d_G(a, b) + d_G(b, x) < d_W(a, b) + d_W(b, x) \leq k/2$ and contradicting that $d_G(a, x) = k/2$. We can now conclude that $d_G(a, b) = d_W(a, b)$ for every a and b in $V(W)$, meaning that W is an isometric cycle of length k . ■

Together the lemmas above yield an equivalence.

Corollary 2.4 *If k is even, there is an isometric cycle of length k in G going through u and v if and only if the distance between (u, v) and (v, u) is $k/2$ in G_k .*

Corollary 2.5 *If k is even, there is an isometric cycle of length k in G if and only if there is a pair of vertices, u and v , with $d_{G_k}[(u, v), (v, u)] = k/2$.*

Now we have an analogous result for odd k 's. For a vertex (u, v) in G_k , we define the set $M'_k(u, v)$ to be the set $\{(u, x) : (u, x) \in V(G_k) \wedge (v, x) \in E(G)\}$. Using this set, we can show the following.

Lemma 2.6 *If k is odd and there is an isometric cycle of length k in G , going through u and v , the distance between (u, v) and a vertex (v, x) in $M'_k(u, v)$ is $\lfloor k/2 \rfloor$ in G_k .*

Proof. Assume G has an isometric cycle of length k , $C = \{c_1, c_2, \dots, c_{\lfloor k/2 \rfloor}, c_{\lfloor k/2 \rfloor + 1}, \dots, c_k\}$ with $c_1 = u$, $c_{\lfloor k/2 \rfloor + 1} = v$ and $c_k = x$. As in the proof of Lemma 2.4, $(c_1, c_{\lfloor k/2 \rfloor + 1})$, $(c_2, c_{\lfloor k/2 \rfloor + 2})$, \dots , $(c_{\lfloor k/2 \rfloor}, c_{k-1})$ and $(c_{\lfloor k/2 \rfloor + 1}, c_k)$ are vertices in G_k . Again we can see that there is an edge between each consecutive pair of vertices, and that $(c_{\lfloor k/2 \rfloor + 1}, c_k)$ must be in $M'_k(v, u)$, because $c_{\lfloor k/2 \rfloor + 1}$ actually is v and there is an edge between c_k and c_1 in G . The given path assures us that $d_{G_k}[(u, v), (v, x)] \leq \lfloor k/2 \rfloor$ while Lemma 2.1 gives $d_{G_k}[(u, v), (v, x)] \geq \lfloor k/2 \rfloor$ and ensures equality. ■

Lemma 2.7 *If k is odd and the distance in G_k between (u, v) and a vertex (v, x) in $M'_k(v, u)$ is $\lfloor k/2 \rfloor$, then there is an isometric cycle of length k in G , going through u and v .*

Proof. Let $d_{G_k}[(u, v), (v, x)] = \lfloor k/2 \rfloor$ and let $P = \{(u, v), (a_2, b_2), (a_3, b_3) \dots (a_{\lfloor k/2 \rfloor}, b_{\lfloor k/2 \rfloor}), (v, x)\}$ be a shortest path between (u, v) and (v, x) . We observe that $W = \{u, a_2, a_3 \dots v, b_2, b_3 \dots, x, u\}$ must be a cyclic walk of length k and a subgraph in G . Assume for a contradiction that there is a pair of vertices a and b in W with $d_G(a, b) < d_W(a, b)$. Let z be a vertex in W so that either (a, z) or (z, a) is in P and $d_W(a, b) + d_W(b, z) \leq \lfloor k/2 \rfloor$. We can always find such a vertex by starting at a and walking $\lfloor k/2 \rfloor$ steps along W either clockwise or counterclockwise. The direction to walk in is the one that makes us visit b on the way to z . But we assumed that $d_G(a, b) < d_W(a, b)$. This means that $d_G(a, z) \leq d_G(a, b) + d_G(b, z) < d_W(a, b) + d_W(b, z) \leq \lfloor k/2 \rfloor$ which obviously contradicts to $d_G(a, x) = \lfloor k/2 \rfloor$. This can only mean that W is an isometric cycle. ■

Corollary 2.8 *If k is odd, there is an isometric cycle of length k in G going through u and v if and only if the distance in G_k between (u, v) and a vertex (v, x) in $M'_k(v, u)$ is $\lfloor k/2 \rfloor$.*

Corollary 2.9 *If k is odd, there is an isometric cycle of length k in G if and only if there are vertices u, v and x so that $x \in M'_k(v, u)$ and $d_{G_k}[(u, v), (v, x)] = \lfloor k/2 \rfloor$.*

In order to simplify notation, and to be able to treat k odd and k even as one case, for a $(u, v) \in V(G_k)$ we define the set $M_k(u, v)$ to be (u, v) if k is even and $M'_k(u, v)$ if k is odd. Now we are able to summarize the above section in one result.

Lemma 2.10 *G has an isometric cycle of length k if and only if there are vertices u and v and $x \in V(G)$ so that $(v, x) \in M_k(v, u)$ and $d_{G_k}[(u, v), (v, x)] = \lfloor k/2 \rfloor$.*

Proof. If k is even the result is equivalent to Corollary 2.5, in the odd case it is the same as Corollary 2.9. ■

3 Algorithm and complexity analysis

It should be clear how one can use G_k to check whether G has an isometric cycle of length k . We use a straightforward approach - For a given k , compute G_k , and find out whether there is a pair of vertices (u, v) and (v, x) in $V(G_k)$ so that $(v, x) \in M_k(v, u)$ and $d_{G_k}[(u, v), (v, x)] = \lfloor k/2 \rfloor$. If such a pair exists we have a cycle of length k , if not - we don't. Now we do this for every k between 3 and n , and this way we find the longest isometric cycle. We observe that Lemma 2.1 guarantees that $d_{G_k}[(u, v), (v, x)] \geq \lfloor k/2 \rfloor$, so we can search for vertices satisfying the inequality $d_{G_k}[(u, v), (v, x)] \leq \lfloor k/2 \rfloor$ instead of the equation above.

Algorithm 3.1**Input:** A graph $G = (V, E)$.**Output:** The length of the longest isometric cycle in G - ans **begin** $ans := 0$ Compute the distance matrix of G .**if** G is a tree **then** $\text{return } ans$ **end-if****for** every k from 3 **to** n **do** $V_k := \emptyset$ **for** every u and v in V **do****if** $d(u, v) = \lfloor k/2 \rfloor$ **then** $V_k := V_k \cup \{(u, v)\}$ **end-if** $E_k := \emptyset$ **for** every (u, v) and (w, x) in V_k **do****if** $(u, w) \in E \wedge (v, x) \in E$ **then** $E_k := E_k \cup \{(u, v), (w, x)\}$ **end-if** $G_k := (V_k, E_k)$ Compute $G_k^{\lfloor k/2 \rfloor} = (V_k, E_k^{\lfloor k/2 \rfloor})$ **for** every triple of vertices (u, v, x) in V **do****if** $(u, v) \in V(G_k) \wedge (v, x) \in M_k(v, u) \wedge [(u, v), (v, x)] \in E_k^{\lfloor k/2 \rfloor}$ **then** $ans := k;$ **end-if****return** ans **end**

Figure 1: The algorithm for computing the longest isometric cycle.

Theorem 3.2 *Given a graph G , Algorithm 3.1 computes the length of the longest isometric cycle of G*

Proof. If G is a tree it has no cycles the algorithm correctly returns 0. If it has a cycle, it must have an isometric cycle of length at least 3 and at most n . Assume the longest isometric cycle of G has length k' . Then Theorem 2.10 states that G_k must have a pair of vertices (u, v) and (v, x) so that $(v, x) \in M(v, u)$ and $d_{G_k}[(u, v), (v, x)] = \lfloor k/2 \rfloor$. This means that $[(u, v), (v, x)]$ must be in $E_k^{\lfloor k/2 \rfloor}$ so the variable ans will be set to k' in the iteration of the outer loop that has $k = k'$. For all iterations after this, e.g. with $k > k'$, the same theorem states that there cannot be any vertices (u, v) and (v, x) that satisfy the above conditions. This ensures that the command $ans := k$ will not be executed. Thus the algorithm terminates with the value of ans equal to k' - the length of G 's longest isometric cycle. ■

Now that we have proven the correctness of Algorithm 3.1 we can move on to analyzing its time complexity and discussing some of the implementation details. As the algorithm is fairly straightforward, the complexity analysis also is quite simple. We can compute the distance matrix and find out whether G is acyclic using naive algorithms in $\mathcal{O}(n^3)$ time. Having pre-computed the distance matrix of G , we now can make queries about the distance between two vertices in G in constant time. Using this we see that V_k is computed in $\mathcal{O}(n^2)$ time for a given k .

Now we observe that E_k is computed in $|V_k|^2$ time. At this point, we arrive at the spot where we have to compute $G_k^{\lfloor k/2 \rfloor}$. The fastest known way to do that is to use the folklore algorithm for computing graph powers; To compute G^x we write x in base 2 and let d_{i+1} be the i 'th digit in this string counting from right to left. Now we find G^{2^k} for $2^k \leq x$ and compute the matrix product $\prod_{i=0}^{\lfloor \log(x) \rfloor} A_i$ where

A_i is defined to be G^{2^i} if $d_i = 1$ and the identity matrix otherwise. It is easy to show that the time complexity of this approach is $|V(G)|^\alpha \log(x)$ when n^α is the time needed to multiply two n by n binary matrices. This means that computing $G_k^{\lfloor k/2 \rfloor}$ from G_k in this manner takes $\mathcal{O}(|V_k|^\alpha \log(\lfloor k/2 \rfloor))$ time.

Having computed $G_k^{\lfloor k/2 \rfloor}$, we only have the last loop left. The loop iterates over all triples (u, v, x) of vertices in V . Having pre-computed G_k , $G_k^{\lfloor k/2 \rfloor}$ and the distance matrix of G , we can perform all the tests in the following if-sentence in $\mathcal{O}(1)$ time. Note that we do not actually compute the set $M_k(v, u)$, we only test whether (v, x) satisfies the conditions to be in the set. To conclude the analysis we just need to summarize the discussion above and make a couple observations.

Observation 3.3 $\sum_{k=3}^n |V_k| \leq 2n^2$

Proof. For given k_1 and k_2 with $\lfloor k_1/2 \rfloor \neq \lfloor k_2/2 \rfloor$ we see that V_{k_1} and V_{k_2} are pairwise disjoint subsets of V^2 . If $\lfloor k_1/2 \rfloor = \lfloor k_2/2 \rfloor$ then $V_{k_1} = V_{k_2}$ by the definition of G_k . By summing over all even and all odd k 's we obtain $\sum_{k=3}^n |V_k| = \sum_{k=1}^{\lfloor (n-1)/2 \rfloor} |V_{2k+1}| + \sum_{k=2}^{\lfloor n/2 \rfloor} |V_{2k}| \leq |V^2| + |V^2| = 2n^2$. ■

Theorem 3.4 *If n^α is the time needed to multiply two n by n binary matrices and $\alpha \geq 2$, Algorithm 3.1 terminates in $\mathcal{O}(n^{2\alpha} \log(n))$ steps.*

Proof. Let T be the total number of steps performed by the algorithm. By the discussion above, we see that $T = \mathcal{O}(n^3) + \sum_{k=3}^n [\mathcal{O}(n^2) + \mathcal{O}(|V_k|^2) + \mathcal{O}(|V_k|^\alpha \log(\lfloor k/2 \rfloor)) + \mathcal{O}(n^3)]$. By rearranging our terms and summing the terms not dependant on k we obtain $T = \mathcal{O}(n^4) + \sum_{k=3}^n \mathcal{O}(|V_k|^\alpha \log(\lfloor k/2 \rfloor))$. Now $\log(\lfloor k/2 \rfloor) = \mathcal{O}(\log(n))$ so we end up with $T = \mathcal{O}(n^4) + \mathcal{O}(\log(n)) \sum_{k=3}^n \mathcal{O}(|V_k|^\alpha)$. As $\alpha \geq 2$, n^α is a convex function and we can put the summation inside the \mathcal{O} . This yields $T = \mathcal{O}(n^4) + \mathcal{O}(\log(n)) \mathcal{O}([\sum_{k=3}^n |V_k|]^\alpha)$ while Observation 3.3 allows us to simplify the expression to $T = \mathcal{O}(n^4) + \mathcal{O}(\log(n)(2n^2)^\alpha)$. As $2\alpha \geq 4$ we may simplify even further, to $T = \mathcal{O}(n^{2\alpha} \log(n))$. ■

Finally, we recall the fastest known algorithm for matrix multiplication.

Theorem 3.5 [1] *Two n by n matrices can be multiplied in $\mathcal{O}(n^{2.376})$*

Corollary 3.6 *Algorithm 3.1 runs in $\mathcal{O}(n^{4.752} \log(n)) = \mathcal{O}(n^{4.753})$ time.*

4 Conclusion

A way to view isometric subgraphs is as a generalization of induced subgraphs. While finding the longest induced cycle in a graph is hard [2], we have seen that finding the longest isometric cycle is easy. Finding the longest isometric path is trivial, as any isometric path must be a shortest path and vice versa. This might lead us to believe that finding out whether a graph G contains a given graph H as an isometric subgraph also might be solvable in polynomial time. However, letting H be the complete graph on k vertices for a given k yields a direct reduction from *Clique* and proves the problem NP-hard. Actually the problem is NP-complete even if we demand that H is a star. To see this, consider the following reduction from Independent Set. For a given graph G construct G' by adding a universal vertex u and connecting u to all vertices of G . It is now easy to see that G' has an isometric star on k edges if and only if G has an independent set of size k .

References

- [1] Don Coppersmith, Shmuel Winograd. Matrix multiplication via arithmetic progressions. *Journal of Symbolic Computation*, v.9 n.3, p.251-280, March 1990
- [2] M. R. Garey and D. S. Johnson. *Computers and Intractability*. W. H. Freeman and Co., 1978.