

Approximating Acyclicity Parameters of Sparse Hypergraphs*

Fedor V. Fomin[†] Petr A. Golovach[†] Dimitrios M. Thilikos[‡]

Abstract

The notions of hypertree width and generalized hypertree width were introduced by Gottlob, Leone, and Scarcello in order to extend the concept of hypergraph acyclicity. These notions were further generalized by Grohe and Marx who introduced the fractional hypertree width of a hypergraph. All these width parameters on hypergraphs are useful for extending tractability of many problems in database theory and artificial intelligence. In this paper, we study the approximability of (generalized, fractional) hypertree width of sparse hypergraphs where the criterion of sparsity reflects the sparsity of their incidence graphs. Our first step is to prove that the (generalized, fractional) hypertree width of a hypergraph \mathcal{H} is constant-factor sandwiched by the treewidth of its incidence graph, when the incidence graph belongs to some apex-minor-free graph class (the family of apex-minor-free graph classes includes planar graphs and graphs of bounded genus). This determines the combinatorial borderline above which the notion of (generalized, fractional) hypertree width becomes essentially more general than treewidth, justifying that way its functionality as a hypergraph acyclicity measure. While for more general sparse families of hypergraphs treewidth of incidence graphs and all hypertree width parameters may differ arbitrarily, there are sparse families where a constant factor approximation algorithm is possible. In particular, we give a constant factor approximation polynomial time algorithm for (generalized, fractional) hypertree width on hypergraphs whose incidence graphs belong to some H -minor-free graph class.

1 Introduction

Many important theoretical and “real-world” problems can be expressed as constrained satisfaction problems (CSP). Among examples one can mention numerous problems from different domains like Boolean satisfiability, temporal reasoning, graph coloring, belief maintenance, machine vision, and scheduling. Another example is the conjunctive-query containment problem, which is a fundamental problem in database query evaluation. In fact, as it was shown by Kolaitis and Vardi [21], CSP, conjunctive-query containment, and finding homomorphism for relational structures are essentially the same problem. The problem is known to be NP-hard in general [3] and polynomial time solvable for restricted class of acyclic queries [29]. Recently, in the database and constraint satisfaction communities various extensions of query

*A preliminary version of this paper appeared in [11].

[†]Department of Informatics, University of Bergen, N-5020 Bergen, Norway. Supported by the Norwegian Research Council.

[‡]Department of Mathematics, University of Athens, Panepistimioupolis, GR15784 Athens, Greece. Supported by the project “Kapodistrias” (AII 02839/28.07.2008) of the National and Kapodistrian University of Athens.

(or hypergraph) acyclicity were studied. The main motivation for the quest for a suitable measure of acyclicity of a hypergraph (query, or relational structure) is the extension of polynomial time solvable cases (like acyclic hypergraphs) to more general instances. In this direction, Chekuri and Rajaraman in [4] introduced the notion of query width. Gottlob, Leone, and Scarcello [14, 15, 17, 18] defined hypertree width and generalized hypertree width. Furthermore, Grohe and Marx [20] have introduced the most general parameter known so far, fractional hypertree width, and proved that CSP, restricted to instances of bounded fractional hypertree width, is polynomial time solvable.

Unfortunately, all known variants of hypertree width are NP-complete [13, 19, 22]. Moreover, generalized hypertree width is NP-complete even when checking whether its value is at most 3 (see [19]). In the case of hypertree width, the problem is $W[2]$ -hard when parameterized by k [13]. Both hypertree width and the generalized hypertree are hard to approximate. For example, the reduction of Gottlob et al. in [13] can be used to show that the generalized hypertree width of an n -vertex hypergraph cannot be approximated within a factor $c \log n$ for some constant $c > 0$ unless $P = NP$.

All these parameters for hypergraphs can be seen as generalizations of the treewidth of a graph. The treewidth is a fundamental graph parameter from Graph Minors Theory by Robertson and Seymour [26] and it has numerous algorithmic applications (for a survey, see [2]). It is an old open question whether the treewidth can be approximated within a constant factor and the best known approximation algorithm for treewidth is $\sqrt{\log OPT}$ -approximation due to Feige et al. [10]. However, as it was shown by Feige et al. [10], the treewidth of an H -minor-free graph is constant factor approximable.

Our results. Our first result is combinatorial. We show that for a wide family of hypergraphs (those where the incidence graph excludes an apex graph as a minor – that is a graph that can become planar after removing a vertex) the fractional and generalized hypertree width of a hypergraph is bounded by a linear function of treewidth of its incidence graph. Apex-minor-free graph classes include planar and bounded genus graphs.

For hypergraphs whose incidence graphs are apex graphs the two parameters may differ arbitrarily, and this result determines the boundary where fractional hypertree width starts being essentially different from treewidth of the incidence graph. This indicates that hypertree width parameters are more useful as the adequate version of acyclicity for non-sparse instances.

Our proof is based on theorems from bidimensionality theory and a min-max (in terms of fractional hyperbrambles) characterization of fractional hypertree width. The proof essentially identifies what is the obstruction analogue of fractional hypertree width for incidence graphs.

Our second result applies further for sparse classes where the difference between (generalized, fractional) hypertree width of a hypergraph and treewidth of its incidence graph can be arbitrarily large. In particular, we give a constant factor approximation algorithm for generalized and fractional hypertree width of hypergraphs with H -minor-free incidence graphs extending the results of Feige et al. [10] from treewidth to (generalized, fractional) hypertree width. The algorithm is based on a series of theorems based on the main decomposition theorem of the Robertson-Seymour's Graph Minor project. As a combinatorial corollary of our results, it follows that generalized hypertree width and fractional hypertree width differ within constant multiplicative factor if the incidence graph of the hypergraph does not contain a fixed graph as a minor.

2 Definitions and preliminaries

2.1 Basic definitions

We consider finite undirected graphs without loops or multiple edges. The vertex set of a graph G is denoted by $V(G)$ and its edge set by $E(G)$ (or simply by V and E if it does not create confusion).

Let G be a graph. For a vertex v , we denote by $N_G(v)$ its (*open*) *neighborhood*, i.e. the set of vertices which are adjacent to v . The *closed neighborhood* of v , i.e. the set $N_G(v) \cup \{v\}$, is denoted by $N_G[v]$. For $U \subseteq V(G)$, we define $N_G[U] = \bigcup_{v \in U} N_G[v]$ (we may omit index if the graph under consideration is clear from the context). If $U \subseteq V(G)$ (or $u \in V(G)$) then $G - U$ (or $G - u$) is the graph obtained from G by the removal of vertices of U (vertex u correspondingly).

Given an edge $e = \{x, y\}$ of a graph G , the graph G/e is obtained from G by contracting e ; which is, to get G/e we identify the vertices x and y and remove all loops and replace all multiple edges by simple edges. A graph H obtained by a sequence of edge-contractions is said to be a *contraction* of G . A graph H is a *minor* of G if H is a subgraph of a contraction of G .

We say that a graph G is *H-minor-free* when it does not contain H as a minor. We also say that a graph class \mathcal{G} is *H-minor-free* (or, excludes H as a minor) when all its members are *H-minor-free*.

An *apex graph* is a graph obtained from a planar graph G by adding a vertex and making it adjacent to some of the vertices of G . A graph class \mathcal{G} is *apex-minor-free* if \mathcal{G} excludes a fixed apex graph H as a minor.

The $(k \times k)$ -*grid* is the Cartesian product of two paths of lengths $k - 1$.

A *surface* Σ is a compact 2-manifold (we always consider connected surfaces). Whenever we refer to a Σ -*embedded graph* G we consider a 2-cell embedding of G in Σ . To simplify notations, we do not distinguish between a vertex of G and the point of Σ used in the drawing to represent the vertex or between an edge and the line representing it. We also consider a graph G embedded in Σ as the union of the points corresponding to its vertices and edges. That way, a subgraph H of G can be seen as a graph H , where $H \subseteq G$. Recall that $\Delta \subseteq \Sigma$ is a (closed) disc if it is homeomorphic to $\{(x, y) : x^2 + y^2 \leq 1\}$. The *Euler genus* of a nonorientable surface Σ is equal to the nonorientable genus $\tilde{g}(\Sigma)$ (or the crosscap number). The *Euler genus* of an orientable surface Σ is $2g(\Sigma)$, where $g(\Sigma)$ is the orientable genus of Σ . We refer to the book of Mohar and Thomassen [24] for more details on graphs embeddings.

If $X \subseteq 2^A$ for some set A , then by $\bigcup X$ we denote the union of all elements of X .

Recall that a *hypergraph* \mathcal{H} is a pair $\mathcal{H} = (V(\mathcal{H}), E(\mathcal{H}))$ where $V(\mathcal{H})$ is a finite nonempty set of vertices, and $E(\mathcal{H})$ is a set of nonempty subsets of $V(\mathcal{H})$ called *hyperedges*, $\bigcup E(\mathcal{H}) = V(\mathcal{H})$. We consider here only hypergraphs without isolated vertices (i.e. every vertex is in some hyperedge).

For vertex $v \in V(\mathcal{H})$, we denote by $E_{\mathcal{H}}(v)$ the set of its incident hyperedges.

The *incidence graph* of the hypergraph \mathcal{H} is the bipartite graph $I(\mathcal{H})$ with vertex set $V(\mathcal{H}) \cup E(\mathcal{H})$ such that $v \in V(\mathcal{H})$ and $e \in E(\mathcal{H})$ are adjacent in $I(\mathcal{H})$ if and only if $v \in e$.

2.2 Treewidth of graphs and hypergraphs

A *tree decomposition* of a hypergraph \mathcal{H} is a pair (T, χ) , where T is a tree and $\chi: V(T) \rightarrow 2^{V(\mathcal{H})}$ is a function associating a set of vertices $\chi(t) \subseteq V(\mathcal{H})$ (called a *bag*) to each node t of the decomposition tree T such that

- i) $V(\mathcal{H}) = \bigcup_{t \in V(T)} \chi(t)$,
- ii) for each $e \in E(\mathcal{H})$, there is a node $t \in V(T)$ such that $e \subseteq \chi(t)$, and
- iii) for each $v \in V(\mathcal{H})$, the set $\{t \in V(T) : v \in \chi(t)\}$ forms a subtree of T .

The *width* of a tree decomposition equals $\max\{|\chi(t)| - 1 : t \in V(T)\}$. The *treewidth* of a hypergraph \mathcal{H} is the minimum width over all tree decompositions of \mathcal{H} . We use notation $\mathbf{tw}(\mathcal{H})$ for the treewidth of a hypergraph \mathcal{H} .

It is easy to verify that for any hypergraph \mathcal{H} , $\mathbf{tw}(\mathcal{H}) + 1 \geq \mathbf{tw}(I(\mathcal{H}))$. However, these parameters can differ considerably on hypergraphs. For example, for the n -vertex hypergraph \mathcal{H} with one hyperedge which contains all vertices, $\mathbf{tw}(\mathcal{H}) = n - 1$ and $\mathbf{tw}(I(\mathcal{H})) = 1$.

Since $\mathbf{tw}(\mathcal{H}) \geq |e|$ for every $e \in E(\mathcal{H})$, we have that the presence of a large hyperedge results in a large treewidth of the hypergraph. The paradigm shift in the transition from treewidth to hypertree width consists in counting the covering hyperedges rather than counting the number of vertices in a bag. This parameter seems to be more appropriate, especially with respect to constraint satisfaction problems. We start with the introduction of even more general parameter of fractional hypertree width.

2.3 Hypertree width, its generalizations and related notions

In general, given a finite set A , we use the term *labelling of A* for any function $\gamma: A \rightarrow [0, 1]$. We also use the notation $\mathcal{G}(A)$ for the collection of all labellings of a set A .

The *size* of a labelling of A is defined as

$$|\gamma| = \sum_{x \in A} \gamma(x).$$

If the values of a labelling γ are restricted to be 0 or 1, then we say that γ is a *binary* labelling of A . Clearly, the size of a binary labelling is equal to the number of the elements of A that are labelled by 1. Given a hyperedge labelling γ of a hypergraph \mathcal{H} , we define the set of vertices of \mathcal{H} that are *blocked* by γ as

$$B(\gamma) = \{v \in V(\mathcal{H}) \mid \sum_{e \in E_{\mathcal{H}}(v)} \gamma(e) \geq 1\},$$

i.e. the set of vertices that are incident to hyperedges whose total labelling sums up to 1 or more.

A *fractional hypertree decomposition* [20] of \mathcal{H} is a triple (T, χ, λ) , where (T, χ) is a tree decomposition of \mathcal{H} and $\lambda: V(T) \rightarrow \mathcal{G}(E(\mathcal{H}))$ is a function, assigning a hyperedge labelling to each node of T , such that for every $t \in V(T)$, $\chi(t) \subseteq B(\lambda(t))$, i.e. all vertices of the bag $\chi(t)$ are blocked by the labelling $\lambda(t)$. The *width* of a fractional hypertree decomposition (T, χ, λ) is $\min\{|\lambda(t)| : t \in V(T)\}$, and the *fractional hypertree width* $\mathbf{fhw}(\mathcal{H})$ of \mathcal{H} is the minimum of the widths of all fractional hypertree decompositions of \mathcal{H} .

If λ assigns a binary hyperedge labeling to each node of T , then (T, χ, λ) is a *generalized hypertree decomposition* [16]. Correspondingly, the *generalized hypertree width* $\mathbf{ghw}(\mathcal{H})$ of \mathcal{H} is the minimum of the widths of all generalized hypertree decompositions of \mathcal{H} .

Clearly, $\mathbf{fhw}(\mathcal{H}) \leq \mathbf{ghw}(\mathcal{H})$ but, as it was shown in [20], there are families of hypergraphs of bounded fractional hypertree width but unbounded generalized hypertree width. Notice that computing the fractional hypertree width is an NP-complete problem even for sparse graphs. To see this, take a connected graph G that is not a tree and construct a new graph H by replacing every edge of G by $|V(G)| + 1$ paths of length 2. It is easy to check that $\mathbf{tw}(G) + 1 = \mathbf{fhw}(H)$ (see also [22]).

The proof of the next lemma follows from results of [4] about query width. For completeness, we provide a direct proof here.

Lemma 1. *For any hypergraph \mathcal{H} , $\mathbf{fhw}(\mathcal{H}) \leq \mathbf{ghw}(\mathcal{H}) \leq \mathbf{tw}(I(\mathcal{H})) + 1$.*

Proof. Let (T, χ) be a tree decomposition of $I(\mathcal{H})$ of width $\leq k$. It is enough to describe a generalized hypertree decomposition (T, χ', λ) for \mathcal{H} that has width $\leq k$. For every $t \in V(T)$, let $\chi'(t) = (\chi(t) - E(\mathcal{H})) \cup (\bigcup(\chi(t) \cap E(\mathcal{H})))$. We include to $\lambda(t)$ all hyperedges $\chi(t) \cap E(\mathcal{H})$, and for every $v \in \chi(t) \cap V(\mathcal{H})$, a hyperedge e such that $v \in e$ is chosen arbitrary and included to $\lambda(t)$. Clearly, $V(\mathcal{H}) = \bigcup_{t \in V(T)} \chi'(t)$, for each $e \in E(\mathcal{H})$ there is a node $t \in V(T)$ such that $e \subseteq \chi'(t)$, and for every $t \in V(T)$ $\chi'(t) \subseteq \bigcup \lambda(t)$. We have to prove that for each $v \in V(\mathcal{H})$, the set $\{t \in V(T) : v \in \chi'(t)\}$ forms a subtree of T . Suppose that there are $s, t \in V(T)$ at distance at least two, $v \in \chi'(s) \cap \chi'(t)$ and $v \notin \chi'(x)$ for all inner vertices x of s, t -path in T . Since (T, χ) is a tree decomposition of $I(\mathcal{H})$, $s \in \chi'(t) - \chi(t)$ or $t \in \chi'(s) - \chi(s)$. Assume that $t \in \chi'(t) - \chi(t)$. It means that there is $e \in \chi(t)$ such that $v \in e$. Note that $e \notin \chi(x)$ for inner vertices x of s, t -path (otherwise $v \in \chi'(x)$ by the definition). If $v \in \chi(s)$ then there is no bag in (T, χ) that contains both endpoints of the edge $\{v, e\} \in E(I(\mathcal{H}))$. So $s \in \chi'(s) - \chi(s)$ and there is $e' \in \chi(s)$ such that $v \in e'$. As before $e' \notin \chi(x)$ for inner vertices and $e \neq e'$. But since v is adjacent with e and e' in $I(\mathcal{H})$, bags $\chi(x)$ contain v and we receive a contradiction. \square

It is necessary to remark here that the fractional hypertree width of a hypergraph can be arbitrarily smaller than the treewidth of its incidence graph. Suppose that a hypergraph \mathcal{H}' is obtained from the hypergraph \mathcal{H} by adding a hyperedge which includes all vertices. Then $\mathbf{fhw}(\mathcal{H}') = 1$ and $\mathbf{tw}(I(\mathcal{H}')) + 1 \geq \mathbf{tw}(I(\mathcal{H})) + 1 \geq \mathbf{fhw}(\mathcal{H})$.

Let \mathcal{H} be a hypergraph. Two sets $X, Y \subseteq V(\mathcal{H})$ *touch* if $X \cap Y \neq \emptyset$ or there exists $e \in E(\mathcal{H})$ such that $e \cap X \neq \emptyset$ and $e \cap Y \neq \emptyset$. A *hyperbramble* of \mathcal{H} is a set \mathcal{B} of pairwise touching connected subsets of $V(\mathcal{H})$ [1]. We say that a labelling γ of $E(\mathcal{H})$ *covers* a vertex set $S \subseteq V(\mathcal{H})$ if some of its vertices are blocked by γ . The *fractional order* of a hyperbramble is the minimum k for which there is a labeling γ of size at most k covering all elements in \mathcal{B} . The *fractional hyperbramble number*, $\mathbf{fhn}(\mathcal{H})$, of \mathcal{H} is the maximum of the fractional orders of all hyperbrambles of \mathcal{H} .

The *robber and army game* was introduced by Grohe and Marx in [20]. The game is played on a hypergraph \mathcal{H} by two players, the robber and the general who commands the army. A position of the game is a pair (γ, v) , where γ is a labelling of $E(\mathcal{H})$ and $v \in V(\mathcal{H})$. The choice of γ is a distribution of the army on the hyperedges of \mathcal{H} , chosen by the general, while v is the position of the robber. During the game, a vertex of the hypergraph is only blocked if the total amount of army on the hyperedges that contain this vertex adds up to the strength of at least one battalion. To start a play of the game, the robber picks a position v_0 , and the initial

position is (\mathcal{O}, v_0) , where \mathcal{O} denote the constant zero mapping. In each round, the players move from the current position (γ, v) to a new position (γ', v') as follows: The general selects γ' , and then the robber selects v' such that there is a path from v to v' in the hypergraph \mathcal{H} that avoids the vertices in $B(\gamma) \cap B(\gamma')$. Under these circumstances, the positions (γ, v) and (γ', v') are called *compatible*. A *game sequence* is a sequence of compatible positions and its cost is the maximum size of a distribution γ in it. If, at some moment, the position of the game is (γ, v) where $v \in B(\gamma)$, then the general wins. If this never happens, then the robber wins. A *winning strategy of cost at most k* for the general is a program that provides a response on each possible position such that any game sequence generated by this program is finite and has cost at most k . The *army width*, $\mathbf{aw}(\mathcal{H})$, of \mathcal{H} is the least k for which there exist a winning strategy of cost at most k .

Using the fact that $\mathbf{aw}(\mathcal{H}) \leq \mathbf{fhw}(\mathcal{H})$ ([20, Theorem 11]), we can prove the following lemma.

Lemma 2. *For any hypergraph \mathcal{H} , $\mathbf{fbn}(\mathcal{H}) \leq \mathbf{fhw}(\mathcal{H})$.*

Proof. Let \mathcal{B} be a hyperbramble of \mathcal{H} of fractional order at least k . Our aim is to provide an escape strategy for the robber against any possible winning strategy of cost at most $< k$. In particular, the robber will always be on a vertex of some set $S \in \mathcal{B}$ such that S not covered by γ and at any position (γ, v) of the game there will be a new unblocked vertex for the robber to move. Indeed, if the response of the general at position (γ, v) is γ' , we have that $|\gamma| < k$ and therefore γ cannot cover all elements of \mathcal{B} . If $S' \in \mathcal{B}$ is such a set, the new position of the robber will be any vertex v' of S' . Clearly, the robber can move from v to v' , as S and S' touch and all of their vertices are unblocked. This implies that $\mathbf{fbn}(\mathcal{H}) \leq \mathbf{aw}(\mathcal{H})$ and the result follows from the fact that $\mathbf{aw}(\mathcal{H}) \leq \mathbf{fhw}(\mathcal{H})$, proved in [20, Theorem 11]. \square

The variant of the robber and army game where the labellings are restricted to be binary labellings is called the *Marshals and Robbers game* and was introduced by Gottlob et al. [18]. The corresponding parameter is called *Marshal width* and is denoted as \mathbf{mw} . Clearly, for any hypergraph \mathcal{H} , $\mathbf{aw}(\mathcal{H}) \leq \mathbf{mw}(\mathcal{H})$.

2.4 i -brambles

An *i -labeled graph* G is a triple (G, N, M) where $N, M \subseteq V(G)$, $N \cup M = V(G)$, $M - N$ and $N - M$ are independent sets of G , and for any $v \in V(G)$ its closed neighborhood $N_G[v]$ is intersecting both N and M . Notice that $\{N, M\}$ is not necessarily a partition of $V(G)$. The incidence graph $I(\mathcal{H})$ of a hypergraph \mathcal{H} can be seen as an i -labeled graph $(I(\mathcal{H}), N, M)$ where $N = V(\mathcal{H})$, $M = E(\mathcal{H})$.

The result of the contraction of an edge $e = \{x, y\}$ of an i -labeled graph (G, N, M) to a vertex v_e is the i -labeled graph (G', N', M') where

- i) $G' = G/e$
- ii) N' contains all vertices of $N - \{x, y\}$ and also the vertex v_e , in case $\{x, y\} \cap N \neq \emptyset$ and
- iii) M' contains all vertices of $M - \{x, y\}$ and also the vertex v_e , in case $\{x, y\} \cap M \neq \emptyset$.

An i -labeled graph (G', N', M') is a *contraction* of an i -labeled graph (G, N, M) if (G', N', M') can be obtained after applying a (possibly empty) sequence of contractions to (G, N, M) . The following lemma is a direct consequence of the definitions.

Lemma 3. *Let (G, N, M) be an i -labeled graph and let G' be a contraction of G . Then there are $N', M' \subseteq V(G')$ such that the i -labeled graph (G', N', M') is a contraction of (G, N, M) .*

Let (G, N, M) be an i -labeled graph. We say that a set $S \subseteq N$ is i -connected if any pair $x, y \in S$ is connected by a path in $G[S \cup M]$. We say that two subsets $S, R \subseteq N$ i -touch if one of the following holds.

- i) $S \cap R \neq \emptyset$, or
- ii) there is an edge $\{x, y\}$ with $x \in S$ and $y \in R$, or
- iii) there is a vertex $z \in M$ such that $N_G[z]$ intersects both S and R .

Given an i -labeled graph (G, N, M) we define an i -bramble of (G, N, M) as any collection \mathcal{B} of i -touching i -connected sets of vertices in N . We say that a labeling γ of M blocks a vertex $x \in N$ if

$$\sum_{y \in N_G[x] \cap M} \gamma(y) \geq 1.$$

We say that γ fractionally covers a vertex set $S \subseteq N$ if some of its vertices is blocked by γ . The order of an i -bramble is the minimum k for which there is a labeling γ of M of size at most k that fractionally covers all sets of \mathcal{B} .

The fractional i -bramble number $\mathbf{fibrn}(G, N, M)$ of an i -labeled graph (G, N, M) is the maximum order of all i -brambles of it.

The following statement follows immediately from the definitions of hyperbrambles and i -brambles.

Lemma 4. *For any hypergraph \mathcal{H} , $\mathbf{fibrn}(I(\mathcal{H}), V(\mathcal{H}), E(\mathcal{H})) = \mathbf{fibrn}(\mathcal{H})$.*

Also it can be easily seen that the fractional i -bramble number is a contraction-closed parameter.

Lemma 5. *If an i -labeled graph (G', N', M') is the contraction of an i -labeled graph (G, N, M) then $\mathbf{fibrn}(G', N', M') \leq \mathbf{fibrn}(G, N, M)$.*

Obviously, i -bramble number is not a subgraph-closed parameter (not even for induced subgraphs), but we can note the following useful claim.

Lemma 6. *Let (G, N, M) be an i -labeled graph and $X \subseteq V(G)$ such that $G - X$ has no isolated vertices, and for every $v \in X \cap M$, $N_G[v] \subseteq X$. Then $(G - X, N - X, M - X)$ is an i -labeled graph and $\mathbf{fibrn}(G - X, N - X, M - X) \leq \mathbf{fibrn}(G, N, M)$.*

Proof. Let $G' = G - X$, $N' = N - X$ and $M' = M - X$. Since G' has no isolated vertices, (G', N', M') is an i -labeled graph. Let \mathcal{B} be an i -bramble of (G', N', M') . Obviously, \mathcal{B} is an i -bramble of (G, N, M) , and there is a labeling γ of M of size $k \leq \mathbf{fibrn}(G, N, M)$ which fractionally covers all sets of \mathcal{B} . It is now enough to note only that the restriction γ' of γ to M is the labeling of M' which covers all sets of \mathcal{B} and $|\gamma'| \leq k$. \square

3 When hypertree width is sandwiched by treewidth

3.1 Influence and valency of i -brambles

Let (G, N, M) be an i -labelled graph and \mathcal{B} an i -bramble of it. We define the *influence* of \mathcal{B} , as $\mathbf{inf}(\mathcal{B}) = \max_{v \in \cup \mathcal{B}} |\{x \in \cup \mathcal{B} \mid \mathbf{dist}_G(v, x) \leq 2\}|$. We also define the *valency* of \mathcal{B} as the quantity $\mathbf{val}(\mathcal{B}) = \max_{v \in \cup \mathcal{B}} |\{S \in \mathcal{B} \mid v \in S\}|$.

Lemma 7. *If \mathcal{B} is an i -bramble of an i -labeled graph (G, N, M) , then the order of \mathcal{B} is at least $\frac{|\mathcal{B}|}{\mathbf{inf}(\mathcal{B}) \cdot \mathbf{val}(\mathcal{B})}$.*

Proof. Let γ be a labelling of M that fractionally covers all sets of \mathcal{B} . We first prove the following claim.

Claim. γ blocks at most $\mathbf{inf}(\mathcal{B}) \cdot |\gamma|$ vertices in $N(\cup \mathcal{B})$.

proof. Let R be a subset of $\cup \mathcal{B}$ such that every vertex in R is blocked by γ . We define G_R as the graph whose vertex set is R and where two vertices $x, y \in R$ are adjacent if their distance in G is 1 or 2. By the definition of influence, we obtain that the maximum degree of G_R is at most $\mathbf{inf}(\mathcal{B}) - 1$ and therefore, G_R has an independent set I of size at least $|R|/\mathbf{inf}(\mathcal{B})$. As $I \subseteq R$, all vertices of I are blocked by γ . This implies that $\forall x \in I \sum_{y \in N_G[x] \cap M} \gamma(y) \geq 1$. By definition, for each pair $x, x' \in I$, $x \neq x'$, $N_G[x] \cap N_G[x'] = \emptyset$. Therefore,

$$|\gamma| = \sum_{x \in M} \gamma(x) \geq \sum_{x \in N_G[R] \cap M} \gamma(x) \geq \sum_{x \in N_G[I] \cap M} \gamma(x) \geq \sum_{x \in I} \sum_{y \in N[x] \cap M} \gamma(y) \geq |I| \geq \frac{|R|}{\mathbf{inf}(\mathcal{B})},$$

and the claim follows.

The above claim, along with the definition of valency, implies that γ fractionally covers no more than $\mathbf{inf}(\mathcal{B}) \cdot |\gamma| \cdot \mathbf{val}(\mathcal{B})$ sets of \mathcal{B} . We conclude that $|\mathcal{B}| \leq \mathbf{inf}(\mathcal{B}) \cdot |\gamma| \cdot \mathbf{val}(\mathcal{B})$ and the lemma follows. \square

3.2 Triangulated grids

A *partially triangulated $(k \times k)$ -grid* is a graph G that is obtained from a $(k \times k)$ -grid (we refer to it as its *underlying grid*) after adding some edges without harming the planarity of the resulting graph. Each vertex of G will be denoted by a pair (i, j) corresponding to its coordinates in the underlying grid. We will also denote as $U(G)$ the vertices, we call them *non-marginal*, of G that in the underlying grid have degree 4 and we call the vertices in $V(G) - U(G)$ *marginal*.

Lemma 8. *Let (G, N, M) be an i -labeled graph, where G is a partially triangulated $(k \times k)$ -grid for $k \geq 4$. Then $\mathbf{fibr}(G, N, M) \geq k/50 - c$, for some constant $c \geq 0$.*

Proof. We use notation $C_{i,j}$ for the set vertices of $N \cap U(G)$ that belong to the i -th row or the j -th column of the underlying grid of G . We claim that $\mathcal{B} = \{C_{i,j} \mid 2 \leq i, j \leq k - 1\}$ is an i -bramble of G of order $\geq k/50 - c$, for some constant $c \geq 0$. Since $k \geq 4$, we have that each set $C_{i,j}$ is non-empty and i -connected. Notice also that the intersection of the i -th row and the j' -th column of the underlying grid of G is either a vertex in N and $C_{i,j} \cap C_{i',j'} \neq \emptyset$, or a vertex in $M - N$, but then all neighbors of it in G belong to N . Therefore, all $C_{i,j}$ and $C_{i',j'}$ should i -touch, and \mathcal{B} is an i -bramble. Each vertex $v = (i, j)$ in $N(\cup \mathcal{B})$ is contained in exactly $2k - 5$ sets of \mathcal{B} (that is $k - 2$ sets $C_{i',j'}$ that agree on the first coordinate plus

$k - 2$ sets $C_{i',j'}$ that agree on the second, minus one set $C_{i,j}$ that agrees on both), therefore $\mathbf{val}(\mathcal{B}) = 2k - 5$. For each non-marginal vertex x in G , there are at most 25 non-marginal vertices within distance ≤ 2 in G (in the worst case, consider a triangulated (5×5) -grid subgraph of G that is centered at x) and thus $\mathbf{ifl}(\mathcal{B}) \leq 25$. As $|\mathcal{B}| = (k - 2)^2$, Lemma 7 implies that there is a constant c such that the order of \mathcal{B} is at least $k/50 - c$ and the lemma follows. \square

We require the following result.

Proposition 1 ([25, (6.2)]). *Let k be a positive integer. Then, every planar graph excluding $(k \times k)$ -grid as a minor has treewidth at most $6k - 5$.*

Theorem 1. *If \mathcal{H} is a hypergraph with a planar incidence graph $I(\mathcal{H})$, then $\mathbf{fhw}(\mathcal{H}) - 1 \leq \mathbf{ghw}(\mathcal{H}) - 1 \leq \mathbf{tw}(I(\mathcal{H})) \leq 300 \cdot \mathbf{fhw}(\mathcal{H}) + c$ for some constant $c \geq 0$.*

Proof. The left hand inequality follows directly from Lemma 1. Suppose now that \mathcal{H} is a hypergraph where $\mathbf{fhw}(\mathcal{H}) \leq k$. By Lemmata 2 and 4, $\mathbf{fhn}(I(\mathcal{H}), V(\mathcal{H}), E(\mathcal{H})) = \mathbf{fhn}(\mathcal{H}) \leq \mathbf{fhw}(\mathcal{H}) \leq k$. By Lemmata 5 and 8, $(I(\mathcal{H}), V(\mathcal{H}), E(\mathcal{H}))$ cannot be i -contracted to an i -labeled graph (G, N, M) where G is a partially triangulated $(l \times l)$ -grid, where $l = 50 \cdot k + O(1)$. By Lemma 3, $\mathcal{I}(\mathcal{H})$ cannot be contracted to a partially triangulated $(l \times l)$ -grid and thus $I(\mathcal{H})$ excludes an $(l \times l)$ -grid as a minor. From Proposition 1, $\mathbf{tw}(I(\mathcal{H})) \leq 6 \cdot l - 5 \leq 300 \cdot k + c$ and the result follows. \square

3.3 Brambles in Gridoids

We call a graph G by a (k, g) -gridoid if it is possible to obtain a partially triangulated $(k \times k)$ -grid after removing at most g edges from it (we call these edges *additional*).

Lemma 9. *Let (G, N, M) be an i -labeled graph where G is a (k, g) -gridoid. Then $\mathbf{fhn}(G, N, M) \geq k/50 - c \cdot g$ for some constant $c \geq 0$.*

Proof. The proof goes the same way as the proof of Lemma 8. The only difference is that now we exclude from \mathcal{B} all the $C_{i,j}$'s where either i or j is the coordinate of some endpoint of an additional edge. Notice that again $\mathbf{val}(\mathcal{B}) \leq 2k - 5$. Moreover, it also holds $\mathbf{ifl}(\mathcal{B}) \leq 25$ as none of the endpoints is in $N(\mathcal{B})$ or $M(\mathcal{B})$. Finally $|\mathcal{B}| \geq (k - 2 - 2 \cdot g)^2$ and the result follows from Lemma 7. \square

We need the following extension of Proposition 1 for graphs of bounded genus.

Proposition 2 ([7, Theorem 4.12]). *Let k be a positive integer. Then, every planar graph excluding $(k \times k)$ -grid as a minor has treewidth at most $6k \cdot (\mathbf{eg}(G) + 1)$.*

The proof of the next theorem is very similar to the one of Theorem 1 (use Lemma 9 instead of Lemma 8 and Proposition 2 instead of Proposition 1).

Theorem 2. *If \mathcal{H} is a hypergraph with an incidence graph $I(\mathcal{H})$ of Euler genus at most g , then $\mathbf{fhw}(\mathcal{H}) - 1 \leq \mathbf{ghw}(\mathcal{H}) - 1 \leq \mathbf{tw}(I(\mathcal{H})) \leq 300 \cdot g \cdot \mathbf{fhw}(\mathcal{H}) + c \cdot g$, for some constant $c \geq 0$.*

3.4 Brambles in augmented grids

An *augmented* $(r \times r)$ -grid of span s is an $r \times r$ grid with some extra edges such that each vertex of the resulting graph is attached to at most s non-marginal vertices of the grid.

Lemma 10. *If (G, N, M) is an i -labeled graph where G is an augmented $(k \times k)$ -grid with span s , then $\mathbf{fibn}(G, N, M) \geq \frac{k}{2 \cdot s^2} - c$, for some constant $c \geq 0$.*

Proof. We consider the i -bramble $\mathcal{B} = \{C_{i,j} \mid 2 \leq i, j \leq k-1\}$ of the proof of Lemma 8 and we directly observe that $\mathbf{val}(\mathcal{B}) \leq 2k - 5$ and $|\mathcal{B}| \geq (k-2)^2$. By the definition of the augmented $(k \times k)$ -grid with span h we obtain that $\mathbf{ifl}(\mathcal{B}) \leq s^2$ and the result follows applying Lemma 7. \square

As it was shown by Demaine et al. [6], every apex-minor-free graph with treewidth at least k can be contracted to a $(f(k) \times f(k))$ -augmented grid of span $O(1)$ (the hidden constants in the “ O ”-notation depend only on the excluded apex). Because, $f(k) = \Omega(k)$ (due to the results of Demaine and Hajiaghayi in [8]), we have the following proposition.

Proposition 3. *Let G be an H -apex-minor-free graph of treewidth at least $c_H \cdot k$. Then G contains as a contraction an augmented $(k \times k)$ -grid of span s_H , where constants c_H, s_H depend only on the size of apex graph H that is excluded.*

The proof of the next theorem is similar to the one of Theorem 1 (use Lemma 10 instead of Lemma 8 and Proposition 3 instead of Proposition 1).

Theorem 3. *If \mathcal{H} is a hypergraph with an incidence graph $I(\mathcal{H})$ that is H -apex-minor-free, then $\mathbf{fhw}(\mathcal{H}) - 1 \leq \mathbf{ghw}(\mathcal{H}) - 1 \leq \mathbf{tw}(I(\mathcal{H})) \leq c_H \cdot \mathbf{fhw}(\mathcal{H})$ for some constant c_H that depends only on H .*

4 Hypergraphs with H -minor-free incidence graphs

The results of Theorem 3 cannot be extended to hypergraphs which incidence graph excludes an arbitrary fixed graph H as a minor. For example, for every integer k , it is possible to construct a hypergraph \mathcal{H} with the planar incidence graph such that $\mathbf{tw}(I(\mathcal{H})) \geq k$. By adding to \mathcal{H} an universal hyperedge containing all vertices of \mathcal{H} , we obtain a hypergraph \mathcal{H}' of generalized hypertree width one. Its incidence graph $I(\mathcal{H}')$ does not contain the complete graph K_6 as a minor, however its treewidth is at least k . Despite of that, in this section we prove that if a hypergraph has H -minor-free incidence graph, then its generalized hypertree width and fractional hypertree width can be approximated by the treewidth of a graph that can be constructed from its incidence graph in polynomial time. By making use of this result we show that in this case generalized hypertree width and fractional hypertree width are up to a constant multiplicative factor from each other. Another consequence of the combinatorial result is that there is a constant factor polynomial time approximation algorithm for both parameters on this class of hypergraphs. Our proof is based on the Excluded Minor Theorem by Robertson and Seymour [27].

4.1 Graph minor theorem

Before describing the Excluded Minor Theorem we need some definitions.

Definition 1 (CLIQUE-SUMS). Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two disjoint graphs, and $k \geq 0$ an integer. For $i = 1, 2$, let $W_i \subseteq V_i$, form a clique of size h and let G'_i be the graph obtained from G_i by removing a set of edges (possibly empty) from the clique $G_i[W_i]$. Let $F : W_1 \rightarrow W_2$ be a bijection between W_1 and W_2 . We define the h -clique-sum of G_1 and G_2 , denoted by $G_1 \oplus_{h,F} G_2$, or simply $G_1 \oplus G_2$ if there is no confusion, as the graph obtained by taking the union of G'_1 and G'_2 by identifying $w \in W_1$ with $F(w) \in W_2$, and by removing all the multiple edges. The image of the vertices of W_1 and W_2 in $G_1 \oplus G_2$ is called the join of the sum.

Note that some edges of G_1 and G_2 are not edges of G , since it is possible that they had edges which were removed by clique-sum operation. Such edges are called *virtual* edges of G . We remark that \oplus is not well defined; different choices of G'_i and the bijection F could give different clique-sums. A sequence of h -clique-sums, not necessarily unique, which result in a graph G , is called a *clique-sum decomposition* of G .

Definition 2 (h -nearly embeddable graphs). Let Σ be a surface with boundary cycles C_1, \dots, C_h , i.e. each cycle C_i is the border of a disc in Σ . A graph G is h -nearly embeddable in Σ , if G has a subset X of size at most h , called apices, such that there are (possibly empty) subgraphs G_0, \dots, G_h of $G - X$ such that

- i) $G - X = G_0 \cup \dots \cup G_h$,
- ii) G_0 is embeddable in Σ , we fix an embedding of G_0 ,
- iii) graphs G_1, \dots, G_h (called vortices) are pairwise disjoint,
- iv) for $1 \leq i \leq h$, let $U_i := \{u_{i_1}, \dots, u_{i_{m_i}}\} = V(G_0) \cap V(G_i)$, G_i has a path decomposition (B_{ij}) , $1 \leq j \leq m_i$, of width at most h such that
 - a) for $1 \leq i \leq h$ and for $1 \leq j \leq m_i$ we have $u_j \in B_{ij}$,
 - b) for $1 \leq i \leq h$, we have $V(G_0) \cap C_i = \{u_{i_1}, \dots, u_{i_{m_i}}\}$ and the points $u_{i_1}, \dots, u_{i_{m_i}}$ appear on C_i in this order (either if we walk clockwise or anti-clockwise).

The following proposition is known as the Excluded Minor Theorem [27] and is the cornerstone of Robertson and Seymour's Graph Minors theory.

Theorem 4 ([27]). For every non-planar graph H , there exists an integer h , depending only on the size of H , such that every graph excluding H as a minor can be obtained by h -clique-sums from graphs that can be h -nearly embedded in a surface Σ in which H cannot be embedded. Moreover, while applying each of the clique sums, at most three vertices from each summand other than apices and vertices in vortices are identified.

Let us remark that by the result of Demaine et al. [9] such a clique-sum decomposition can be obtained in time $O(n^c)$ for some constant c which depends only from H (see also [5]).

4.2 Approximation

Let \mathcal{H} be a hypergraph such that its incidence graph $G = I(\mathcal{H})$ excludes a fixed graph H as a minor. Every graph excluding a planar graph H as a minor has a constant treewidth [25]. Thus if H is planar, by the results of Theorem 3, the generalized hypertree width does not exceed some constant. In what follows, we always assume that H is not planar.

By Theorem 4, there is an h -clique-sum decomposition of $G = G_1 \oplus G_2 \oplus \dots \oplus G_m$ such that for every $i \in \{1, 2, \dots, m\}$, the summand G_i can be h -nearly embedded in a surface Σ in which H can not be embedded. We assume that this clique-sum decomposition is *minimal*, in the sense that for every virtual edge $\{x, y\} \in E(G_i)$ there is an x, y -path in G with all inner vertices in $V(G) - V(G_i)$ (otherwise it is always possible to remove such edges and modify clique-sum operations correspondingly). Let A_i be the set of apices of G_i . We define $E_i = A_i \cap E(\mathcal{H})$ and $G'_i = G_i - (N_G[E_i] \cup A_i)$. For every virtual edge $\{x, y\}$ of G'_i we perform the following operation: if there is no x, y -path in $G - (N[E_i] \cup A_i)$ with all inner vertices in $G - V(G'_i)$, then $\{x, y\}$ is removed from G'_i . We denote the resulted graph by F_i .

In what remains we show that the maximal value of $\mathbf{tw}(F_i)$, where maximum is taken over all $i \in \{1, 2, \dots, m\}$, is a constant factor approximation of generalized and fractional hypertree widths of \mathcal{H} . The upper bound is given by the following lemma. Its proof uses the fact that $\mathbf{ghw}(\mathcal{H}) \leq 3 \cdot \mathbf{mw}(\mathcal{H}) + 1$ (see [1]) and is based on the description of a winning strategy for $k = \max\{\mathbf{tw}(F_i) : i \in \{1, 2, \dots, m\}\} + 2h + 1$ marshals on \mathcal{H} .

Lemma 11. $\mathbf{ghw}(\mathcal{H}) \leq 3 \cdot \max\{\mathbf{tw}(F_i) : i \in \{1, 2, \dots, m\}\} + 6h + 4$.

Proof. Let $w = \max\{\mathbf{tw}(F_i) : i \in \{1, 2, \dots, m\}\}$ and $k = w + 2h + 1$. By the result of Adler et al. [1], we have that $\mathbf{ghw}(\mathcal{H}) \leq 3 \cdot \mathbf{mw}(\mathcal{H}) + 1$, and it is enough to describe a winning strategy for k marshals on \mathcal{H} .

The clique-sum decomposition $G = G_1 \oplus G_2 \oplus \dots \oplus G_m$ can be considered as a tree decomposition (T, χ) of G for some tree T with nodes $\{1, 2, \dots, m\}$ such that $\chi(i) = V(G_i)$, i.e. the vertex sets of the summands are the bags of this decomposition. The idea behind the winning strategy for marshals is to “chase” the robber in the hypergraph along $m + 1$ decompositions for its incidence graph: one is induced by the clique-sum decomposition and others are tree decompositions of F_i . We say that marshals *block* a set $X \subseteq V(G)$ if all hyperedges $X \cap E(\mathcal{H})$ are occupied by them, and for every $v \in X \cap V(\mathcal{H})$, there is a hyperedge $e \in E(\mathcal{H})$, occupied by a marshal, such that $v \in e$.

Let us note that the definition of F_i yields the following: if $x, y \in V(F_i)$ and there is a x, y -path in $G - (N[E_i] \cup A_i)$ with all inner vertices not in F_i , then $\{x, y\}$ is an edge of F_i . (Indeed, if $\{x, y\}$ is an edge of G , then it is also an edge of F_i . If $\{x, y\} \notin E(G)$ but such a path exists, then $\{x, y\}$ is a virtual edge in G_i and by the definition of F_i , such an edge is also an edge of F_i .)

For $i \in \{1, 2, \dots, m\}$, let $(T^{(i)}, \chi_i)$ be a tree decomposition of F_i of width at most w . We assume that trees T and $T^{(1)}, T^{(2)}, \dots, T^{(m)}$ are rooted trees with roots r and r_1, r_2, \dots, r_m respectively.

For a node $i \in V(T)$ and its parent j (in T), we define $S = V(G_i) \cap V(G_j)$. (If $i = r$ then we put $S = \emptyset$.) By the definition of the clique-sum, $|S| \leq h$. Assume that at most h marshals are already placed on the hypergraph in such a way that they block S . Assume also that the robber occupies some vertex of $\chi(T^{(i)})$. We put at most h marshals on hyperedges to block the set of apices A_i . Then the set $N_G[E_i] \cup A_i$ is blocked by these marshals.

Now marshals start to “chase” the robber in the subhypergraph induced by the vertex set $V(F_i) \cap V(\mathcal{H})$ along $T^{(i)}$. We put at most $w + 1$ marshals to block the set $\chi_i(r_i)$. Assume now that some set $\chi_i(x)$ for $x \in V(T^{(i)})$ is blocked, and that the robber can only occupy vertices of $\chi_i(T_y^{(i)})$, where $T_y^{(i)}$ is a subtree of $T^{(i)}$ rooted in some child y of the node x . We remove some marshals which were placed to block $\chi_i(x)$ in such a way that $\chi_i(x) \cap \chi_i(y)$ remains blocked, and then place additional marshals to block $\chi_i(y)$. This manoeuvre can be done by

making use of at most $w + 1$ marshals. We put $x = y$ and repeat this operation until there is a child y of x such that the robber can be in $\chi_i(T_y^{(i)})$. Thus by repeating at most $|V(T^{(i)})|$ times this operation, marshals “push” the robber out of $V(F_i) \cap V(\mathcal{H})$.

Let j be a child of i in T such that the robber now can occupy only the vertices of $\chi(T_j)$, where T_j is the subtree of T rooted at j . Let $S' = V(G_i) \cap V(G_j)$. Since $|S'| \leq h$, we have that h marshals can block this set and, after that, all other marshals can be removed from \mathcal{H} .

We apply the described strategy of marshals starting from $i = r$ until the robber is captured in some leaf-node of T . For every node of T we have used at most h marshals to occupy apices, at most h marshals to block the vertices of the clique-sum, and at most $w + 1$ marshals to push the robber out of F_i . Thus in total at most $2h + w + 1$ marshals have a winning strategy on \mathcal{H} . \square

We also need a result roughly stating that if a graph G with a big grid as a surface minor is embedded on a surface Σ of small genus, then there is a disc in Σ containing a big part of the grid of G . This result is implicit in the work of Robertson and Seymour and there are simpler alternative proofs by Mohar and Thomassen [23, 28] (see also [7, Lemma 3.3]). We use the following variant of this result from Geelen et al. [12].

Proposition 4 ([12]). *Let g, l, r be positive integers such that $r \geq g(l + 1)$ and let G be an (r, r) -grid. If G is embedded in a surface Σ of Euler genus at most $g^2 - 1$, then some (l, l) -subgrid of G is embedded in a closed disc Δ in Σ such that the boundary cycle of the (l, l) -grid is the boundary of the disc.*

Now we are ready to prove the following lower bound.

Lemma 12. $\text{fbn}(\mathcal{H}) \geq \varepsilon_H \cdot \max\{\text{tw}(F_i) : i \in \{1, 2, \dots, m\}\}$ for some constant ε_H depending only on H .

Proof. Let $i \in \{1, 2, \dots, m\}$. We assume that $G - (N[E_i] \cup A_i)$ is a connected graph which has at least one edge. (Otherwise one can consider the components of this graph separately and remove isolated vertices.) The main idea of the proof is to contract it to a planar graph with approximately the same treewidth as F_i and then apply same techniques that were used in the previous section for the planar case.

Structure of $G - (N[E_i] \cup A_i)$. Let us note that an h -clique-sum decomposition $G = G_1 \oplus G_2 \oplus \dots \oplus G_m$ induces an h -clique-sum decomposition of $G' = G - (N[E_i] \cup A_i)$ with the summand G_i replaced by F_i . Let G'_1, G'_2, \dots, G'_l be the connected components of $G' - V(F_i)$. Every such component G'_j is attached via clique-sum to F_i by some clique Q_j of F_i . Note that cliques Q_j contain all virtual edges of F_i . We assume that each clique Q_j does not separate vertices of F_i . Otherwise, it is possible to decompose F_i into the clique-sum of graphs $F_i^{(1)} \oplus F_i^{(2)}$ with the join Q_j and prove the bound for summands and, since $\text{tw}(F_i) = \max\{F_i^{(1)}, F_i^{(2)}\}$, that will prove the lemma. To simplify the structure of the graph, for every component G'_j , we contract all its edges and denote by S_j the star whose central vertex is the result of the contraction and leaves are the vertices of Q_j .

Contracting vortices. The h -nearly embedding of the graph G_i induces the h -nearly embedding of $F_i = X_0 \cup X_1 \cup \dots \cup X_h$ without apices. Here we assume that X_0 is embedded in a surface Σ of genus depending on H and X_1, X_2, \dots, X_h are the vortices. For every vortex X_j ,

the vertices $V(X_0) \cap V(X_j)$ are on the boundary C_j of some face of X_0 . If for a star S_k some of its leaves Q_k are in X_j or C_j , we do the following operation: if $Q_k \cap (V(X_j) - V(C_j)) \neq \emptyset$ then all edges of S_k are contracted, and if $Q_k \cap (V(X_j) - V(C_j)) = \emptyset$ but $|Q_k \cap V(C_j)| \geq 2$, then we contract all edges of S_k that are incident to the vertices of $Q_k \cap V(C_j)$. These contractions result to the contraction of some edges of F_i . Particularly, all virtual edges of X_j and C_j are contracted. Additionally, we contract all remaining edges of X_j and C_j . We perform these contractions for all vortices of F_i and denote the result by F'_i . It follows immediately from the definition of the h -clique-sum and the fact that at most three vertices that do not belong in vortices or apices are identified, that F'_i coincides with the graph obtained from F_i by contracting all vortices X_j and all boundaries of faces C_j . It can be easily seen that F'_i is embedded in Σ . It is known (see e.g. [7,8]) that there is a positive constant a_H which depends only on H such that $\mathbf{tw}(F'_i) \geq a_H \cdot \mathbf{tw}(F_i)$.

Contracting the part that lies outside of some planar disc. Since F'_i is embedded in Σ , we have that the graph F'_i contains some $(k \times k)$ -grid as a surface minor, where $k \geq b_H \cdot \mathbf{tw}(F'_i)$ for some constant b_H [7]. Combining this result with Proposition 4, we obtain the following claim.

Claim. There is a disc $\Delta \subseteq \Sigma$ such that

- i) the subgraph R of F'_i induced by vertices of $F'_i \cap \Delta$ is a connected graph;
- ii) the subgraph R' of F'_i induced by $N_{F'_i}[V(R)]$ is completely in some disc Δ' ;
- iii) vertices of $V(R') - V(R)$ induce a cycle C which is the border of Δ' , and
- iv) $\mathbf{tw}(R) \geq c_H \cdot \mathbf{tw}(F'_i)$ for some constant c_H .

The claim above permits us to treat the part of F'_i which is outside Δ exactly in the same way we have treated vortices. For stars S_k intersecting $V(F'_i) - V(R')$ or C , we do the following: if $Q_k \cap (V(F'_i) - V(R')) \neq \emptyset$, then all edges of S_k are contracted, and if $Q_k \cap (V(F'_i) - V(R')) = \emptyset$ but $|Q_k \cap V(C)| \geq 2$, then all edges of S_k incident to the vertices of $Q_k \cap V(C)$ are contracted. These contractions result in the contraction of some edges of F'_i with endpoints on C or outside Δ' . Particularly, all such virtual edges are contracted. Additionally, we contract all remaining edges of $F'_i - V(R)$ and C . Thus this part of the graph is contracted to a single vertex. Denote the obtained graph X . This graph is planar, and since R is a subgraph of X , we have that $\mathbf{tw}(X) \geq \mathbf{tw}(R)$.

Embedding the stars. Some edges of X are virtual, and all such edges are in cliques Q_j . By the fact that while taking clique sums at most three vertices that do not belong in vortices or apices are identified, we obtain that $|Q_j| \leq 3$. For every clique $Q = V(X) \cap Q_j$, we do the following. If $Q = \{x, y\}$, then the edge of the star S_j incident to x is contracted. If $Q = \{x, y, z\}$, then if two vertices of Q , say x and y , are joined by an edge in G , then the edge of S_j incident to z is contracted, and if there are no such edges and the triangle induced by $\{x, y, z\}$ is the boundary of some face of X , then we add a new vertex on this face, join it with x, y and z (it can be seen as S_j embedded in this face, and since our graph is i -labeled, it is assumed that this new vertex has same labels as the central vertex of S_j), and then remove virtual edges. Note that if the triangle is not a boundary of some face, then Q is a separator of our graph, but we assumed that there are no such separators. Denote by Y the obtained

graph. Similar construction was used in the proof of the main theorem in [8] (Theorem 1.2), and by the same arguments as were used by Demaine et al. we immediately conclude that there is a positive constant d_H such that $\mathbf{tw}(X) \geq d_H \cdot \mathbf{tw}(Y)$.

Now all contractions are finished. Note that the graph Y is a planar graph which is a contraction of $G' = G - (N[E_i] \cup A_i)$. Also there is some positive constant e_H which depends only on H such that $\mathbf{tw}(Y) \geq e_H \cdot \mathbf{tw}(F_i)$. Recall that we consider the i -labeled graph $(G, V(\mathcal{H}), E(\mathcal{H}))$. By Lemma 4, $\mathbf{fhn}(\mathcal{H}) = \mathbf{fhn}(G, V(\mathcal{H}), E(\mathcal{H}))$. Because the sets $V(\mathcal{H})$ and $E(\mathcal{H})$ are independent, by Lemma 6, we have that $\mathbf{fhn}(G, V(\mathcal{H}), E(\mathcal{H})) \geq \mathbf{fhn}(G', N, M)$, where $N = V(\mathcal{H}) - (N[E_i] \cup A_i)$ and $M = E(\mathcal{H}) - (N[E_i] \cup A_i)$. By Lemma 5, $\mathbf{fhn}(G', N, M) \geq \mathbf{fhn}(Y, N', M')$, where N' and M' are sets which were obtained as the result of contractions of N and M . Finally, as in Theorem 1, one can show that $\mathbf{fhn}(Y, N', M') \geq f_H \cdot \mathbf{tw}(Y)$ for some constant f_H . By putting all these bounds together, we prove that there is a positive constant ε_H which depends only on H , such that $\mathbf{fhn}(\mathcal{H}) \geq \varepsilon_H \cdot \mathbf{tw}(F_i)$. \square

Combining Lemmata 1, 2, 11, and 12, we obtain the following theorem.

Theorem 5. $(1/c_H) \cdot w \leq \mathbf{fhw}(\mathcal{H}) \leq \mathbf{ghw}(\mathcal{H}) \leq c_H \cdot w$, where $w = \max\{\mathbf{tw}(F_i) : i \in \{1, 2, \dots, m\}\}$, and c_H is a constant depending only on H .

Remark. Notice that, by Theorem 5, the generalized hypertree width and the fractional hypertree width of a hypergraph with H -minor-free incidence graph may differ within a multiplicative constant factor. We stress that, as observed in [20], this is not the case for general hypergraphs.

Demaine et al. [9] (see also [5, 10, 27]) described an algorithm which constructs a clique-sum decomposition of an H -minor-free graph G on n vertices with the running time $n^{O(1)}$ (the hidden constant in the running time depends only on H). As far as we constructed summands G_i , the construction of graphs F_i can be done in polynomial time. Moreover, since the algorithm of Demaine et al. provides h -nearly embeddings of these graphs, it is possible to use it to construct a polynomial constant factor approximation algorithm for the computation of $\mathbf{tw}(F_i)$. This provides us with the main algorithmic result of this section.

Theorem 6. *For any fixed graph H , there is a polynomial time c_H -approximation algorithm computing the generalized hypertree width and the fractional hypertree width for hypergraphs with H -minor-free incidence graphs, where the constant c_H depends only on H .*

5 Conclusion

Let us remark that while the winning strategy for marshals used in the proof of Lemma 11 is not monotone (a strategy is *monotone* if the territory available for the robber only decreases in the game), but it can be turned into monotone by choosing marshals' positions in a slightly more careful way. By making use of the results from [18], the monotone strategy can be used to construct a generalized hypertree decomposition (or fractional hypertree decomposition). Thus our results can be used not only to approximate but to construct the corresponding decompositions as well.

A long-standing open in Graph Algorithms is whether the treewidth of a planar graph is computable in polynomial time. It would be very interesting to see, if an NP-hardness proof can be obtained for at least one of the hypertree width parameters of planar hypergraphs. Moreover, a polynomial time algorithm for any of these hypertree width parameters on planar

hypergraphs would be a significant breakthrough. While we do not know if there exists a polynomial time algorithm for any of these problems, it would be challenging to ask if some of the variants of the problem is fixed parameter tractable on planar hypergraphs. On the other hand, the treewidth of a planar graphs admits a constant factor approximation. Our results extend this algorithmic result to all the hypertree width parameters on planar hypergraphs.

Finally, the sparsity of hypergraphs studied in this paper is expressed in terms of their incidence graphs. It is an interesting question whether there are other sparsity measures where further algorithmic or complexity results can be obtained for hypertree width parameters.

References

- [1] I. ADLER, G. GOTTLOB, AND M. GROHE, *Hypertree width and related hypergraph invariants*, European J. Combin., 28 (2007), pp. 2167–2181.
- [2] H. L. BODLAENDER, *A partial k -arboretum of graphs with bounded treewidth*, Theoret. Comput. Sci., 209 (1998), pp. 1–45.
- [3] A. K. CHANDRA AND P. M. MERLIN, *Optimal implementation of conjunctive queries in relational data bases*, in STOC’77, ACM, 1977, pp. 77–90.
- [4] C. CHEKURI AND A. RAJARAMAN, *Conjunctive query containment revisited*, Theoret. Comput. Sci., 239 (2000), pp. 211–229.
- [5] A. DAWAR, M. GROHE, AND S. KREUTZER, *Locally excluding a minor*, in LICS’07, IEEE Computer Society, 2007, pp. 270–279.
- [6] E. D. DEMAINE, F. V. FOMIN, M. HAJIAGHAYI, AND D. M. THILIKOS, *Bidimensional parameters and local treewidth*, SIAM J. Discrete Math., 18 (2004/05), pp. 501–511.
- [7] ———, *Subexponential parameterized algorithms on graphs of bounded genus and H -minor-free graphs*, J. ACM, 52 (2005), pp. 866–893.
- [8] E. D. DEMAINE AND M. HAJIAGHAYI, *Graphs excluding a fixed minor have grids as large as treewidth, with combinatorial and algorithmic applications through bidimensionality*, in SODA’05, ACM, 2005, pp. 682–689.
- [9] E. D. DEMAINE, M. T. HAJIAGHAYI, AND K. ICHI KAWARABAYASHI, *Algorithmic graph minor theory: Decomposition, approximation, and coloring*, in FOCS’05, IEEE Computer Society, 2005, pp. 637–646.
- [10] U. FEIGE, M. HAJIAGHAYI, AND J. R. LEE, *Improved approximation algorithms for minimum weight vertex separators*, SIAM J. Computing, 38 (2008), pp. 629–657.
- [11] F. V. FOMIN, P. A. GOLOVACH, AND D. M. THILIKOS, *Approximating acyclicity parameters of sparse hypergraphs*, in 26th Symposium on Theoretical Aspects of Computer Science (STACS 2009), 2009.
- [12] J. F. GEELLEN, R. B. RICHTER, AND G. SALAZAR, *Embedding grids in surfaces*, European J. Combin., 25 (2004), pp. 785–792.

- [13] G. GOTTLOB, M. GROHE, N. MUSLIU, M. SAMER, AND F. SCARCELLO, *Hypertree decompositions: Structure, algorithms, and applications*, in WG'05, vol. 3787 of Lecture Notes in Computer Science, Springer, 2005, pp. 1–15.
- [14] G. GOTTLOB, N. LEONE, AND F. SCARCELLO, *A comparison of structural CSP decomposition methods*, Artificial Intelligence, 124 (2000), pp. 243–282.
- [15] ———, *The complexity of acyclic conjunctive queries*, J. ACM, 48 (2001), pp. 431–498.
- [16] ———, *Hypertree decompositions and tractable queries*, J. Comput. System Sci., 64 (2002), pp. 579–627.
- [17] ———, *Robbers, marshals, and guards: game theoretic and logical characterizations of hypertree width*, J. Comput. System Sci., 66 (2003), pp. 775–808.
- [18] ———, *Robbers, marshals, and guards: game theoretic and logical characterizations of hypertree width*, J. Comput. System Sci., 66 (2003), pp. 775–808.
- [19] G. GOTTLOB, Z. MIKLÓS, AND T. SCHWENTICK, *Generalized hypertree decompositions: NP-hardness and tractable variants*, in PODS'07, ACM, 2007, pp. 13–22.
- [20] M. GROHE AND D. MARX, *Constraint solving via fractional edge covers*, in SODA'06, ACM, 2006, pp. 289–298.
- [21] P. G. KOLAITIS AND M. Y. VARDI, *Conjunctive-query containment and constraint satisfaction*, J. Comput. System Sci., 61 (2000), pp. 302–332.
- [22] D. MARX, *Approximating fractional hypertree width*, in SODA'09, ACM, 2009.
- [23] B. MOHAR, *Combinatorial local planarity and the width of graph embeddings*, Canad. J. Math., 44 (1992), pp. 1272–1288.
- [24] B. MOHAR AND C. THOMASSEN, *Graphs on surfaces*, Johns Hopkins University Press, Baltimore, MD, 2001.
- [25] N. ROBERTSON, P. SEYMOUR, AND R. THOMAS, *Quickly excluding a planar graph*, J. Combin. Theory Ser. B, 62 (1994), pp. 323–348.
- [26] N. ROBERTSON AND P. D. SEYMOUR, *Graph minors. II. Algorithmic aspects of tree-width*, J. Algorithms, 7 (1986), pp. 309–322.
- [27] ———, *Graph minors. XVI. Excluding a non-planar graph*, J. Combin. Theory Ser. B, 89 (2003), pp. 43–76.
- [28] C. THOMASSEN, *A simpler proof of the excluded minor theorem for higher surfaces*, J. Combin. Theory Ser. B, 70 (1997), pp. 306–311.
- [29] M. YANNAKAKIS, *Algorithms for acyclic database schemes*, in VLDB'81, IEEE Computer Society, 1981, pp. 82–94.