

Upper bounds on boolean-width with applications to exact algorithms

Yuri Rabinovich¹, Jan Arne Telle², and Martin Vatshelle²

¹ Department of Computer Science, University of Haifa, Israel

² Department of Informatics, University of Bergen, Norway

Abstract. Boolean-width is similar to clique-width, rank-width and NLC-width in that all these graph parameters are constantly bounded on the same classes of graphs. In many classes where these parameters are not constantly bounded, boolean-width is distinguished by its much lower value, such as in permutation graphs and interval graphs where boolean-width was shown to be $O(\log n)$ [1]. Together with FPT algorithms having runtime $O^*(c^{boolw})$ for a constant c this helped explain why a variety of problems could be solved in polynomial-time on these graph classes.

In this paper we continue this line of research and establish non-trivial upper-bounds on the boolean-width and linear boolean-width of *any* graph. Again we combine these bounds with FPT algorithms having runtime $O^*(c^{boolw})$, now to give a common framework of moderately-exponential exact algorithms that beat brute-force search for several independence and domination-type problems, on general graphs.

Boolean-width is closely related to the number of maximal independent sets in bipartite graphs. Our main result breaking the triviality bound of $n/3$ for boolean-width and $n/2$ for linear boolean-width is proved by new techniques for bounding the number of maximal independent sets in bipartite graphs.

1 Introduction

Boolean-width is a recently introduced graph parameter motivated by algorithms [2]. Having small boolean-width is witnessed by a decomposition of the graph into cuts with few different unions of neighborhoods - Boolean sums of neighborhoods - across the cut. This makes the decomposition natural to guide dynamic programming algorithms to solve problems where vertex sets having the same neighborhood across a cut can be treated as equivalent. Such dynamic programming on a given decomposition of boolean-width $boolw$ will for several problems related to independence and domination have runtime $O^*(c^{boolw})$ for a small constant c [2].

Boolean-width is similar to clique-width, rank-width and NLC-width in that all these graph parameters are constantly bounded on the same classes of graphs. However, in many classes where these parameters are not constantly bounded, boolean-width is distinguished by its much lower value. For example, permutation graphs, interval graphs, convex graphs and Dilworth k graphs all have boolean-width $O(\log n)$, and the decompositions are easy to find [1]. Since $O^*(c^{O(\log n)})$ is $n^{O(1)}$ this helps explain why several problems related to independence and domination are polynomial-time solvable on these graph classes.

In this paper we continue this line of research, combining $O^*(c^{boolw})$ dynamic programming for independence and domination problems with new bounds on boolean-width. Rather than giving a framework for polynomial-time algorithms on restricted graph classes, our goal in this paper is a framework for moderately-exponential exact algorithms on general graphs. Our main results are non-trivial upper-bounds of $(1-c)n/3$ on the boolean-width, and $(1-c)n/2$ on the linear boolean-width, of *any* graph, for some $c > 0$ and sufficiently large values of n . This is accompanied by a polynomial-time algorithm computing a decomposition witnessing the non-trivial bound on linear boolean-width.

We combine this with dynamic programming algorithms on decompositions of linear boolean-width k that solve INDEPENDENT SET in time $O^*(2^k)$ and DOMINATING SET, INDEPENDENT DOMINATING SET and TOTAL DOMINATING SET in time $O^*(2^{2k})$. The combination gives moderately-exponential exact algorithms on general graphs solving all these problems, also weighted versions and counting versions, by a runtime beating brute-force search. Note that faster algorithms do exist in the literature, our goal in this paper is mainly to show the viability of this line of research. This is the first time a non-trivial upper bound on the value of a graph width parameter has been shown to hold *for every graph*.

Boolean-width is defined based on branch decompositions of a graph, using as cut function what is called the boolean dimension $\text{bd}(H)$ of a (bipartite) graph H . This corresponds to the logarithm (base 2) of the number of maximal independent sets $\text{mis}(H)$ of H . Our upper bounds on (linear) boolean-width rely on new techniques for bounding the number of maximal independent sets. This number has received much attention both from the algorithmic and the structural perspectives. While it is known [10] that computing $\text{mis}(G)$ is $\#P$ -hard even for planar bipartite graphs, approximating it is a much more delicate problem. From the structural point of view, bounding the number of maximal independent sets in special as well as in general graphs leads to interesting hard problems. Let us just mention the entropy-based results about $\text{bd}(G) = \log_2 \text{mis}(G)$ of d -regular graphs [8], see the references therein for an updated picture of the state of research in this area.

We introduce three techniques for our bounds. The first (Theorem 5) is based on a vertex partition achieved from a packing of paths and goes via $\text{im}(H)$, the size of a maximum induced matching in bipartite graph H . The second (Theorem 6) is based on a random partition and also goes via $\text{im}(H)$. The third (Theorem 7) is based on Hoeffding's inequality and in contrast to the first two applies also to boolean-width rather than just linear boolean-width. As already mentioned, our goal is to show the viability of this line of research and these various techniques should be helpful in later attempts to improve the bounds.

Our paper is organized as follows. In Section 2 we give all definitions and some preliminary results, for example showing that a non-trivial upper bound on (linear) boolean-width of a graph G will follow from a non-trivial upper bound on the boolean dimension of some balanced partition of G . In Section 3 we aim at an understanding of the structure of bipartite graphs of high boolean dimension. It is well known and easy that $\text{bd}(H)$ is at most $n/2$, and the maximum is attained by a size- $n/2$ matching. We tie $\text{bd}(H)$ to $\text{im}(H)$ and introduce and study the values of $\text{co-im}(H) = n/2 - \text{im}(H)$

and $\text{co-bd}(H) = n/2 - \text{bd}(H)$, the high-end ranges of $\text{im}(H)$ and $\text{bd}(H)$. We show constant factor approximation algorithms for these values, as well as a stability result showing that the smaller the values are, the closer is the bipartite graph to the size- $n/2$ matching. In Section 4 we turn to general graphs, and show, constructively, by a polynomial-time algorithm, that every graph has a balanced partition where the boolean dimension of the associated bipartite graph beats the triviality bound. Combined with the result from Section 2 this implies a constructive result for linear boolean-width of general graphs, beating the triviality bound of $n/2$. In Section 5 we turn to the standard boolean-width parameter and show also in this case a non-trivial upper bound beating the triviality bound of $n/3$, also constructive, but now by a randomized and low-exponential-time algorithm.

2 Terminology and Preliminaries

We consider undirected unweighted simple graphs $G = (V, E)$ and bipartite graphs $H = (A, B, E)$. We also denote the vertex set V by $V(G)$. For $S \subseteq V$ we denote by $G|_S$ the subgraph induced by S . The *neighborhood* of a vertex $v \in V$ is denoted $N(v)$. The neighborhood of a set $S \subset V$ is $N(S) = \cup_{v \in S} N(v)$. Any $S \subseteq V$ defines a cut $(S, V-S)$, and a bipartite graph $G_{S, V-S} = (S, V-S, \{(u, v) \in E : u \in S \wedge v \in V-S\})$.

A *decomposition tree* of a graph $G = (V, E)$ is a pair (T, δ) where T is a ternary tree, i.e. all internal nodes are of degree three, and δ a bijection between the leaves of T and $V(G)$. Removing an edge (a, b) from T results in two subtrees T_a and T_b , and a bipartition of V into V_a and V_b corresponding, respectively, to the δ -labels of leaves of T_a and T_b , and a bipartite graph G_{V_a, V_b} .

Definition 1 (Boolean Dimension, Boolean Width and Linear Boolean Width). For a bipartite graph $H = (A, B, E)$, let $\mathcal{N}_A = \{N(X) \subseteq B \mid X \subseteq A\}$ be the family of neighborhoods of all sets $X \subseteq A$. The boolean dimension of H is defined as $\text{bd}(H) = \log_2 |\mathcal{N}_A|$.

The boolean-width of a decomposition tree (T, δ) is the maximum value of $\text{bd}(G_{V_a, V_b})$ over all edges (a, b) of T . The boolean-width of G , denoted $\text{bw}(G)$, is the minimum boolean-width over all decomposition trees of G .

The linear boolean-width of G , denoted $\text{lbw}(G)$, is the minimum boolean-width over all decomposition trees (T, δ) of G where T is a path on $|V|$ inner nodes, each with an attached leaf, corresponding to a linear arrangement of V .

Given a graph G there is a $O^*(2.52^n)$ algorithm computing its boolean-width exactly [11] and in FPT time parameterized by $\text{bw}(G)$ we can compute a decomposition of boolean-width $2^{2\text{bw}(G)}$ using the algorithm for decompositions of optimal rank-width [6]. The boolean-width parameter was originally introduced in [2] in the context of parameterized algorithms. In particular, using a natural dynamic programming approach it was shown there that

Theorem 1. [2, 11] Given a graph G and a decomposition tree of boolean-width k , one can solve weighted and counting versions of INDEPENDENT SET in time $O^*(2^{2k})$ and DOMINATING SET, INDEPENDENT DOMINATING SET and TOTAL DOMINATING SET in time $O^*(2^{3k})$.

These are dynamic programming algorithms that choose a root of the decomposition tree and traverse it bottom-up, with each node of the tree representing the subgraph induced by vertices corresponding to the leaves of the subtree. In a decomposition tree for linear boolean-width we choose one end of the path of inner nodes as root so that one of the two children of any node will always represent a subgraph on a single vertex. The runtime on linear decompositions can for this reason be improved

Corollary 1. [2, 11] *Given a linear arrangement of $V(G)$ of linear boolean-width k , one can solve weighted and counting versions of INDEPENDENT SET in time $O^*(2^k)$ and DOMINATING SET, INDEPENDENT DOMINATING SET and TOTAL DOMINATING SET in time $O^*(2^{2k})$.*

Note that for any bipartite graph $H = (A, B, E)$, we have $|\mathcal{N}_A| = |\mathcal{N}_B|$, see, e.g., [9]. A good combinatorial way to demonstrate this is by establishing a bijection between the elements of \mathcal{N}_A (or \mathcal{N}_B) and the set of all maximal independent sets of $H = (A, B, E)$. Here is a sketch of the argument. Given a set $S \in \mathcal{N}_A$, let X be the maximal set in A such that $N(X) = S$. Then X is uniquely defined, and $X \cup B - S$ is maximal independent. In the other direction, given a maximal independent set I , $B - I \in \mathcal{N}_A$. Moreover, if I resulted from S , then S results from I .³ Hence,

Proposition 1. *Let $\text{mis}(H)$ be the number of maximal independent sets in a bipartite graph H . Then, $\text{bd}(H) = \log_2 \text{mis}(H)$.*

The following simple property of $\text{bd}(H)$ will prove useful; it is an immediate consequence of the definition of bd .

Proposition 2. *$\text{bd}(H)$ is monotone decreasing with respect to vertex removal. Moreover, such removal may decrease $\text{bd}(H)$ by at most 1. Hence, for a bipartite graph $H = (A, B, E)$ we have $\text{bd}(H) \leq \min(|A|, |B|)$.*

More generally, given two bipartite graphs $G = (A, B, E)$ and $H = (A, B, E')$ on the same vertex set and the same two sides, it holds that $\text{bd}(G \cup H) \leq \text{bd}(G) + \text{bd}(H)$.

Proposition 2 implies that $\text{bd}(H) \leq n/2$, and this bound is met when H is a matching of size $n/2$. For a finer study of the structure of sub-extremal graphs H , we shall need the following notions.

Definition 2. *Define $\text{im}(G)$ as the size of a maximum induced matching in G , i.e. a maximum-size set of edges whose endpoints do not induce any other edges in G . Note that $\text{im}(G) \leq n/2$ and this bound is met only by a size- $n/2$ matching. To study the high-end range of $\text{bd}(H)$ and $\text{im}(G)$ we define $\text{co-im}(G) = \frac{n}{2} - \text{im}(G)$ and the boolean co-dimension $\text{co-bd}(H) = \frac{n}{2} - \text{bd}(H)$.*

The extremal values of $\text{bw}(G)$ and $\text{lbw}(G)$ are not known. The following proposition provides a preliminary tool for the study of the former.

³ Observed by Nathann Cohen in a course of discussion with the authors. Later we have learned that a similar observation was made in [3].

Proposition 3. *Let $A \subseteq V$ be a subset of vertices with $\frac{1}{3}n \leq |A| \leq \frac{2}{3}n$ and $\text{bd}(G_{A,V-A}) = (\frac{1}{3} - \epsilon)n$ for some $\frac{1}{4} > \epsilon \geq 0$. Then one can construct a decomposition tree of boolean-width at most $(\frac{1}{3} - \frac{\epsilon}{3})n$.*

In particular, $\text{bw}(G) \leq n/3$. We call this the triviality bound for $\text{bw}(G)$.

Proof. Without loss of generality, $|A| \leq n/2$; otherwise we switch to $V-A$. Partition V into three sets $A \cup X, B_1, B_2$ where X is disjoint from A and $|X| \leq \frac{2}{3}\epsilon n$, $|A \cup X| \leq n/2$ and $|B_1|, |B_2| \leq (\frac{1}{3} - \frac{1}{3}\epsilon)n$. Refining this partition to a decomposition tree arbitrarily with the sole restriction that at the upper level $A \cup X$ is split into two approximately equal parts we argue that the proposition holds using Proposition 2. For the top cut $\text{bd}(G_{A \cup X, V-(A \cup X)}) \leq n/3 - \epsilon + |X| \leq (\frac{1}{3} - \frac{\epsilon}{3})n$. For any node representing a subset of B_1 or B_2 the proposition holds since the size of one side will be small. When splitting $A \cup X$ into two approximately equal parts, each part has size at most $n/4$, we have $n/4 \leq n/3 - \epsilon/3$ since $\epsilon < n/4$, and hence the proposition holds. The conclusion about $\text{bw}(G) \leq n/3$ corresponds to choosing an A of size $\frac{1}{3}n$, and $\epsilon = 0$. \square

For $\text{lbw}(G)$ one has an analogous statement:

Proposition 4. *Let $A \subseteq V$ be subset of vertices of size $n/2$ such that $\text{bd}(G_{A,V-A}) \leq (\frac{1}{2} - \epsilon) \cdot n$ for some $\epsilon \geq 0$. Then one can construct a linear arrangement of the vertices of linear boolean-width at most $(\frac{1}{2} - \frac{1}{2}\epsilon) \cdot n$.*

In particular, $\text{lbw}(G) \leq n/2$. We call this the triviality bound for $\text{lbw}(G)$.

Proof. Take any linear arrangement whose first $n/2$ elements are precisely A . We claim that it has the desired property. Let A_i denote the set of the first (equivalently, the last) i elements in this arrangement. Let $G_i = G_{A_i, V-A_i}$. Proposition 2 implies that $\text{bd}(G_i) \leq i$. It also implies that $|\text{bd}(G_i) - \text{bd}(G_{i+1})| \leq 1$ for every i , hence $\text{bd}(G_i) \leq \text{bd}(G_{n/2}) + (n/2 - i) \leq n - \epsilon n - i$. Combining the two bounds on $\text{bd}(G_i)$, the statement follows. \square

The present paper, besides studying $\text{bd}(H)$, is mostly dedicated to establishing upper bounds on $\text{lbw}(G)$ and $\text{bw}(G)$. Before starting with our toil, let us just mention that the above values can in general be as large as $\Omega(n)$, which is achieved e.g., when G is a constant-degree expander. Indeed, in this case any $G_{A,V-A}$ where both sides are $\geq n/3$ has at least $\Omega(n)$ edges, and hence, due to the constant degree, $\text{im}(G_{A,V-A}) = \Omega(n)$. Since the size of an induced matching is a lower bound on the boolean dimension (see the next section for details), the conclusion follows. The constants obtained along this line of reasoning are, however, quite miserable. The extremal values of $\text{lbw}(G)$, $\text{bw}(G)$ and the structure of the corresponding extremal graphs remain a (highly inspiring) mystery.

3 On Boolean co-Dimension

3.1 Boolean Dimension vs. the size of Maximum Induced Matching

We start with a lemma relating the boolean dimension of a bipartite graph $G = (A, B, E)$, $|V(G)| = n$, to the size of the maximum induced matching in G :

Lemma 1. *When $\text{im}(G) \leq n/4$, it holds that*

$$\text{im}(G) \leq \text{bd}(G) \leq \text{im}(G) \cdot \log_2(n/\text{im}(G)) \cdot \phi(2 \cdot \text{im}(G)/n),$$

where ϕ is a function which never exceeds 1.088, and tends to 1 as $\text{im}(G)$ tends to $n/4$.

Proof. The first inequality is obvious, as the boolean dimension is monotone with respect to taking induced subgraphs. For the second inequality, assume w.l.o.g., that $|A| \leq n/2$, and consider the family of neighborhoods \mathcal{N}_A in B . For every $S \in \mathcal{N}_A$, there is a minimal set $S^* \subseteq A$ such that $N(S^*) = S$. By minimality of S^* , each vertex v^* in it has a neighbour $v \in S$ not seen by the other vertices. Forming a set $S' \subseteq S \subseteq B$ by picking (one) such v for every $v^* \in S^*$, we conclude that the subgraph of G induced by (S^*, S') is an induced matching. In particular, it holds that $|S^*| \leq \text{im}(G)$, and thus any $S \in \mathcal{N}_A$ is a neighbourhood of a subset of A of size $\leq \text{im}(G)$. Consequently, using a standard estimation for the sum of binomial coefficients,

$$2^{\text{bd}(G)} = |\mathcal{N}_A| \leq \sum_{i=0}^{\text{im}(G)} \binom{|A|}{i} \leq \sum_{i=0}^{\text{im}(G)} \binom{n/2}{i} \leq 2^{n/2 \cdot H(\text{im}(G)/(n/2))}$$

where $H(p) = p \log_2 \frac{1}{p} + (1-p) \log_2 \frac{1}{1-p}$ is the entropy function. We introduce $p \log_2 \frac{1}{p} + p = p \log_2 \frac{2}{p}$ as an approximator of H , and set $\phi(p) = H(p)/(p \log_2 \frac{2}{p})$. Then,

$$\text{bd}(G) \leq \frac{n}{2} \cdot H\left(\frac{\text{im}(G)}{n/2}\right) = \text{im}(G) \cdot \log_2\left(\frac{n}{\text{im}(G)}\right) \cdot \phi\left(\frac{\text{im}(G)}{n/2}\right)$$

Through numerical analysis we found that $\phi(0.157) \approx 1.08798$ is the global maximum of ϕ in the range $[0, 0.5]$. \square

Thus, $\text{im}(G)$ is a log n -approximation of $\text{bd}(G)$, and the quality of approximation improves as $\text{im}(G)$ grows. However, when $\text{im}(G) \geq n/4$, the approach of Lemma 1 fails to imply anything beyond the trivial upper bound $\text{bd}(G) \leq n/2$. This makes Lemma 1 inapplicable to the study of $\text{co-bd}(G)$ vs. $\text{co-im}(G)$. The key result of this subsection is that when $\text{co-bd}(G)$ is small, so is $\text{co-im}(G)$, and, moreover, $\text{co-bd}(G)$ and $\text{co-im}(G)$ are linearly related.

We start with a special case:

Lemma 2. *Let $G = (A, B, E)$ be a bipartite graph of degree at most 2. Then,*

$$\text{co-im}(G) \geq \text{co-bd}(G) \geq 0.339 \cdot \text{co-im}(G).$$

Proof. The first inequality was already established in Lemma 1. For the second inequality, observe that both $\text{co-bd}(G)$ and $\text{co-im}(G)$ are additive with respect to disjoint union of graphs. Thus it suffices to consider connected G 's, i.e. G is either C_n , the (even) n -cycle, or P_{n-1} , the path on $(n-1)$ edges. For such graphs both $\text{im}(G)$ and $\text{bd}(G)$ are tractable. For the maximum induced matching one easily gets $\text{im}(C_n) = \lfloor \frac{n}{3} \rfloor$ and $\text{im}(P_{n-1}) = \lfloor \frac{n+1}{3} \rfloor$. For boolean dimension, recall that by Proposition 1, $\text{bd}(G) = \log_2 \text{mis}(G)$, where $\text{mis}(G)$ is the number of maximal independent sets in

G . Let $c(n) = \text{mis}(C_n)$ and $p(n) = \text{mis}(P_{n-1})$. The recurrence formulae for these values are well known (see e.g. [5]). Namely, $c(n) = c(n-2) + c(n-3)$, and $p(n) = p(n-2) + p(n-3)$. The initial conditions are $c(1) = 0$, $c(2) = 2$, $c(3) = 3$ and $p(1) = 1$, $p(2) = 2$, $p(3) = 2$ respectively.

Thus, to compare $\text{co-bd}(G)$ to $\text{co-im}(G)$ one needs to lower-bound the expressions

$$\frac{n/2 - \log_2(c(n))}{n/2 - \lfloor \frac{n}{3} \rfloor} \quad \text{and} \quad \frac{n/2 - \log_2(p(n))}{n/2 - \lfloor \frac{n+1}{3} \rfloor}.$$

Combining case analysis (according to $n \bmod 3$), numerical computations and an inductive argument, we conclude that the minimum is achieved on the 8-cycle C_8 , and its value is $\frac{1}{2}(4 - \log_2 10) \approx 0.339036$. \square

We continue with the general case.

Theorem 2. *Let $G = (A, B, E)$ be a bipartite graph. Then,*

$$\text{co-im}(G) \geq \text{co-bd}(G) \geq 0.0698 \cdot \text{co-im}(G) - 4.$$

Proof. As before, we shall be concerned only with the second inequality. Set $\Delta = \text{co-bd}(G)$. Keeping in mind that $\text{mis}(G) = 2^{\text{bd}(G)}$, observe that

$$\text{mis}(G) \leq \text{mis}(G|_{V-\{v\}}) + \text{mis}(G|_{V-\{v\}-N(v)}), \quad (1)$$

where the first term (over-)counts the maximal independent sets not containing v , and the second term counts those containing v . In accordance with inequality, we define a splitting process, or a (weighted) rooted splitting tree T , as follows.

Each inner node x of T is labelled by (G_x, v) , where G_x is an induced subgraph of G , and $v \in V(G_x)$ is a vertex of degree 3 or more in G_x . At the root $G_x = G$; the leaves correspond to induced subgraphs of degree at most 2. An inner node x has two children, one corresponding to the graph obtained from G_x by removing v , the other corresponding to the graph obtained from G_x by removing v and all its neighbours (at least 4 vertices removed). The weight of the respective edge is defined as the number of vertices (respectively) removed. The weight of the node x , $w(x)$, is defined as the sum of weights on the path from the root to x .

In view of (1), it holds that

$$2^{\text{bd}(G)} \leq \sum_{x: \text{leaf of } T} 2^{\text{bd}(G_x)}. \quad (2)$$

The strategy of proof is as follows. The leaves L of T will be split into $L^+ = \{x \mid w(x) \geq z\}$ and $L^- = \{x \mid w(x) < z\}$ according to a suitably defined threshold value z . Then, it is shown that the leaves in L^+ contribute little to the above sum, while the graphs G_x corresponding to $x \in L^-$ have co-im comparable with Δ . The value of z will be set later, in the course of analysis.

Upper-bounding the contribution of L^+

By Proposition 2, $\text{bd}(G_x) \leq (n - w(x))/2$ for any $x \in T$. Thus,

$$\sum_{x \in L^+} 2^{\text{bd}(G_x)} \leq 2^{n/2} \cdot \sum_{x \in L^+} 2^{-w(x)/2}.$$

It is readily checked that the right-hand side is maximized when T is the complete 1-4 tree, where every inner node has an outgoing edge of weight 1 and an outgoing edge of weight 4. Moreover, for an inner node x and its two children x_1 and x_2 it holds that

$$2^{-w(x_1)/2} + 2^{-w(x_2)/2} = 2^{-w(x)/2} \cdot (2^{-1/2} + 2^{-4/2}) < 2^{-w(x)/2}.$$

Therefore, the right-hand side is maximized when the leaves are immediate descendants of the inner nodes of weight $< z$.

Let $s(i)$ denote the number of nodes of weight i in the complete 1-4 tree. Then, $|L^+| \leq s(z) + s(z+1) + s(z+2) + s(z+3)$. A closer look reveals that $s(0), s(1), s(2) = 1$, $s(3) = 2$, and that for $i \geq 4$, $s(i) = s(i-1) + s(i-4)$, where the first term counts the strings with leading "1", and the second term counts those with leading "4". Moreover, it holds that $s(z) + s(z+1) + s(z+2) + s(z+3) = s(z+6)$. Finding the maximal absolute-value root $\alpha = 1.38028$ of the equation $x^4 = x^3 + 1$, and using the estimation $s(z) \leq \alpha^z$, we conclude that

$$|L^+| \leq s(z+6) \leq 8 \cdot 1.38028^z \leq 8 \cdot 2^{0.4649589z},$$

and the total contribution of L^+ to the right-hand side of (2) is bounded from above by

$$2^{n/2} \cdot \sum_{x \in L^+} 2^{-w(x)/2} \leq 8 \cdot 2^{n/2} \cdot 2^{-z/2} \cdot 2^{0.4649589z} \leq 8 \cdot 2^{n/2} \cdot 2^{-0.035z}.$$

Setting $z = \lceil (\Delta + 4)/0.035 \rceil$, where $\Delta = \text{co-bd}(G) = n/2 - \text{bd}(G)$, ensures that the total contribution of L^+ is at most $0.5 \cdot 2^{n/2} \cdot 2^{-\Delta} = 0.5 \cdot 2^{\text{bd}(G)}$.

Consequently, the total contribution of L^- to the right-hand side of (2) is at least

$$\sum_{x \in L^-} 2^{\text{bd}(G_x)} \geq 0.5 \cdot 2^{\text{bd}(G)} = 0.5 \cdot 2^{n/2} \cdot 2^{-\Delta-1}. \quad (3)$$

Upper bounding the contribution of L^-

To get an estimation from above on $\sum_{x \in L^-} 2^{\text{bd}(G_x)}$, consider $\text{bd}(G_x)$ for a leaf x of T . Since G_x has degree ≤ 2 , Lemma 2 implies that:

$$\text{bd}(G_x) = (n - w(x))/2 - \text{co-bd}(G_x) \leq (n - w(x))/2 - 0.339 \cdot \text{co-im}(G_x).$$

Keeping in mind that $w(x) = n - |V(G_x)|$, it follows that $\text{co-im}(G_x) \geq \text{co-im}(G) - w(x)/2$. Substituting this in the previous line yields

$$\text{bd}(G_x) \leq n/2 - 0.33w(x) - 0.339 \cdot \text{co-im}(G),$$

and thus,

$$\sum_{x \in L^-} 2^{\text{bd}(G_x)} \leq \sum_{x \in L^-} 2^{\frac{n}{2} - 0.33w(x) - 0.339\text{co-im}(G)} = 2^{\frac{n}{2} - 0.339\text{co-im}(G)} \sum_{x \in L^-} 2^{-0.33w(x)}$$

As before, the complete 1-4 tree yields the most general (i.e., the weakest possible) upper bound on the sum $\sum_{x \in L^-} 2^{-0.33w(x)}$, as it maximizes the number of nodes of any weight i in T . Since this time the contribution of the father node is dominated by

that of its sons, it suffices to analyse the case when the leaves of L^- have weights $z - 4, z - 3, z - 2$ or $z - 1$. Arguing as before, we conclude that $|L^-| \leq s(z + 2) \leq 2^{0.4649589(z+2)} < 2^{0.4649589z+1}$. That is,

$$\sum_{x \in L^-} 2^{-0.33 w(x)} \leq |L^-| \cdot 2^{-0.33(z-4)} \leq 2^{0.4649589z+1} \cdot 2^{-0.33(z-4)} < 2^{0.1349589z+2.32}.$$

Now, $z < (\Delta + 4)/0.035 + 1$, implying $0.1349589z + 2.32 \leq 3.856\Delta + 18$. The bottom line is:

$$\sum_{x \in L^-} 2^{\text{bd}(G_x)} \leq 2^{n/2} \cdot 2^{-0.339 \cdot \text{co-im}(G)} \cdot 2^{3.856\Delta+18}. \quad (4)$$

We are ready to conclude the proof of Theorem 2. Combining (3) and (4) yields

$$2^{n/2} \cdot 2^{-\Delta-1} \leq \sum_{x \in L^-} 2^{\text{bd}(G_x)} \leq 2^{n/2} \cdot 2^{-0.339 \cdot \text{co-im}(G)} \cdot 2^{3.856\Delta+18}.$$

Combining the two sides, it follows that $0.339 \cdot \text{co-im}(G) \leq 4.856\Delta + 19$, and, finally, $\text{co-im}(G) \leq 14.33(\Delta + 4) = 14.33(\text{co-bd}(G) + 4)$. \square

One curious structural implication following at once from Theorem 2 is the following result (for asymptotically tight results see [8]):

Corollary 2. *Let G be a d -regular bipartite graph, $d > 1$. Then $\text{mis}(G) \leq 2^{(\frac{1}{2}-\epsilon)n}$ for some universal $\epsilon > 0$.*

The reason is that by a trivial computation, for such graphs one has $\text{im}(G) \leq \frac{n}{2} \cdot \frac{d}{2d-1}$, and since $\text{co-bd}(G)$ is proportional to $\text{co-im}(G)$, the conclusion follows.

3.2 A Constant Factor Polynomial Approximation Algorithm for $\text{co-bd}(G)$

Let us first give a polynomial time constant-factor approximation algorithm for $\text{co-im}(G)$. As before, G is bipartite.

Approx-CoIm: *Construct (greedily or otherwise) a maximal vertex-disjoint packing \mathcal{P} of P_2 's (paths on 2 edges) in G . Remove all the vertices in \mathcal{P} . Output \widetilde{M} , the set of the remaining edges.*

Theorem 3. *The above algorithm produces an induced matching \widetilde{M} with $n/2 - |\widetilde{M}| \leq 5 \cdot \text{co-im}(G)$. In particular, it provides a 5-approximation for $\text{co-im}(G)$.*

Proof. Observe that after the removal of P_2 's in \mathcal{P} , the remaining induced graph consists of singletons and isolated edges, and thus \widetilde{M} is indeed an induced matching.

Let M^* denote the maximum induced matching of G . Since every P_2 in the packing must contain at least one vertex outside of M^* , the size of \mathcal{P} is at most $n - 2|M^*|$. Now, since each P_2 in \mathcal{P} may hit at most 2 edges of M^* , at least $|M^*| - 2|\mathcal{P}|$ edges of M^* will survive the removal of \mathcal{P} . Thus,

$$\begin{aligned} |\widetilde{M}| &\geq |M^*| - 2|\mathcal{P}| \geq |M^*| - 2 \cdot (n - 2|M^*|) = 5|M^*| - 2n; \\ n/2 - |\widetilde{M}| &\leq 5/2 n - 5|M^*| = 5 \cdot (n/2 - |M^*|) = 5 \cdot \text{co-im}(G) \square. \end{aligned}$$

Theorem 3 combined with Theorem 2 yields a constant factor approximation algorithm for $\text{co-bd}(G)$:

Theorem 4. *The co-size of \widetilde{M} produced by **Appox-CoIm** on input G , i.e., $n/2 - |\widetilde{M}|$, approximates $\text{co-bd}(G)$ within a multiplicative factor of $5 \cdot 14.3 < 72$.*

4 Linear Boolean Width: Beyond the Triviality Bound

We show (constructively and efficiently) that every size- n graph has a balanced bipartition of its vertex set such that the boolean dimension of the associated bipartite graph is $\leq (1/2 - c)n$ for some universal $c > 0$. Combined with Proposition 4, this implies that $\text{lbw}(G) \leq (1/2 - c/2)n$; the argument therein provides also the corresponding linear arrangement of the vertices.

Two constructions are provided, the first deterministic and somewhat elaborate, the other is just the random uniform bipartition. We start with the former.

GoodBipartition: *Construct (greedily or otherwise) a maximal vertex-disjoint packing \mathcal{P} of P_2 's (paths on 2 edges) in G . For each $P_2 \in \mathcal{P}$ mark the middle vertex. Partition the vertices into two equal- (up to ± 2) size sets so that*

- (i) *for every $P_2 \in \mathcal{P}$, the marked and the unmarked vertices lay on different sides;*
- (ii) *no edge (parity permitting) remaining after the removal of $V(\mathcal{P})$ is split.*

Since \mathcal{P} is maximal, removing $V(\mathcal{P})$ one obtains an (induced) graph that consists of isolated edges and vertices. The required partition is obtained by placing the middle vertices of P_2 's, one by one, on alternating sides. The same is done for surviving isolated edges and isolated vertices.

Let $H = (A, B, E')$ be the graph defined by this bipartition.

Theorem 5. *It holds that $\text{co-im}(H) \geq \frac{1}{10}n$, and $\text{bd}(H) \leq (\frac{1}{2} - \frac{1}{143})n + O(1)$.*

Proof. The edge set of H consists of all the edges of \mathcal{P} , plus some edges of \mathcal{C} , the cut defined by $V(\mathcal{P})$. To bound $\text{im}(H)$, consider $\text{im}(H^*)$ where $H^* = (V, E(\mathcal{P}) \cup \mathcal{C})$. Clearly, $\text{im}(H) \leq \text{im}(H^*)$. Let M^* be the maximum induced matching of H^* .

Each $P_2 \in \mathcal{P}$ is adjacent to some $e \in M^*$, since otherwise M^* could be extended. If some $P_2 \in \mathcal{P}$ is adjacent to a single edge $e \in M^*$, replace (if needed) e in M^* by an adjacent edge in $E(P_2)$, obtaining another maximum induced matching of H^* . Let $H^{**} = (V, M^* \cup E(\mathcal{P}))$. By above observation, H^{**} is w.l.o.g., a disjoint union of $|\mathcal{P}|$ simple paths or/and cycles of (vertex) size ≥ 3 . Consequently, any connected component of H^{**} of size k contributes to M^* at most

$$\text{im}(C_k) \leq \text{im}(P_{k-1}) = \left\lfloor \frac{k+1}{3} \right\rfloor \leq \frac{2}{5}k$$

edges, with equality achieved (only) at P_4 . Thus, $|M^*| \leq \frac{2}{5}n$. Since $\text{im}(H) = |M^*|$, we conclude that $\text{im}(H) \leq \frac{2}{5}n$, and, equivalently, $\text{co-im}(H) \geq \frac{1}{2}n - \frac{2}{5}n = \frac{1}{10}n$. By Theorem 2, this implies that $\text{co-bd}(H) \geq \frac{1}{10 \cdot 14.3}n = \frac{1}{143}n$. \square

Next, we show that a statement similar to that of Theorem 5 holds also for a random uniform bipartition of $V(G)$, when each vertex of V is assigned randomly and independently either to side A or to side B , resulting in $H(A, B, E'')$. While the bound will be weaker, this structural result is of independent interest. The proof is in the appendix.

Theorem 6. *For H as above, it holds almost surely that $\text{co-im}(H) \geq \frac{1}{28}n - o(n)$. Consequently, $\text{co-bd}(H) \geq \frac{1}{801}n - o(n)$.*

5 Boolean Width: Beyond the Triviality Bound

We turn to the boolean-width of general graphs. Since we currently have much less understanding of boolean dimension of unbalanced partitions than of the balanced ones, in this section we provide an existential argument, which can nevertheless be turned into an exponential time algorithm with a relatively small exponent.

Lemma 3. *Every graph G has $A \subset V(G)$ with $\frac{n}{3} \leq |A| \leq \frac{n}{2}$ such that $\text{bd}(G_{A, V-A}) \leq \frac{n}{3} - \frac{n}{226} + o(n)$.*

Proof. If there exists $S \subseteq V(G)$ such that $|S| = \frac{n}{226}$ and $|N(S) \cup S| \leq \frac{n}{3}$ we just take a set A of size $n/3$ containing $N(S) \cup S$. Then, $\text{bd}(G_{A, V-A}) \leq \frac{n}{3} - \frac{n}{226}$ since no vertex in S has a neighbour in $V-A$.

Otherwise, every set S of size $\frac{n}{226}$ has $|N(S) - S| \geq \frac{n}{3} - \frac{n}{226}$. The set A will be constructed by a random procedure by choosing every vertex v with probability $1/3$, randomly and independently from the others. We claim that almost surely the two events hold: first, $|A| = (1 - o(1)) \cdot n/3$, and second, *all* sets $S \subseteq V-A$ of size $\frac{n}{226}$ have $|N(S) \cap A| > (\frac{1}{3} - 0.202) \cdot (\frac{n}{3} - \frac{n}{226})$, which we short-cut as αn for a suitable α . Such an A will be called good.

Since $1 - \Pr(X \cap Y) \leq (1 - \Pr(X)) + (1 - \Pr(Y))$, it suffices to show that each of the two events holds almost surely *separately*. To bound the probabilities of failure, we use the Hoeffding Bound. Let X be the number of successes in r i.i.d. 0/1 events, each happening with probability p . Then by [7],

$$\Pr(X \leq (p - t)r) \leq e^{-2t^2 r}.$$

The desired bound on the probability of the first event follows at once. For the second event, the analysis is more technical.

Let S be any subset of size $\frac{n}{226}$. The probability that S causes a failure is

$$\begin{aligned} & \Pr(\{S \subset V-A\} \wedge \{|N(S) \cap A| \leq \alpha n\}) = \\ & = \Pr(\{|N(S) \cap A| \leq \alpha n\} \mid \{S \subset V-A\}) \cdot \Pr(S \subset V-A). \end{aligned}$$

The probability of the first factor in the above product can be bounded as follows. The set S is fixed and is in $V-A$. The choosing process on the remaining $V-S$ vertices remains, however, unaltered. Since the vertices in $N(S) \setminus S$ are chosen randomly and independently as before, and there are $n/3 - n/226$ vertices in this set, taking $t = 0.202$, we get from Hoeffdings bound:

$$\Pr\left(|N(S) \cap A| \leq \left(\frac{1}{3} - 0.202\right) \left(\frac{n}{3} - \frac{n}{226}\right) \mid \{S \subset V-A\}\right) <$$

$$< e^{-2 \cdot 0.202^2 (n/3 - n/226)} < e^{-0.0268n}.$$

We now use the union bound summing over all sets S of size $\frac{n}{226}$.

$$\begin{aligned} \Pr(\text{There exists a bad } S) &\leq e^{-0.0268n} \cdot \Pr(S \subset V-A) \cdot \binom{n}{n/226} = \\ &= e^{-0.0268n} \cdot \binom{2n/3}{n/226} \approx e^{-0.0268n} \cdot 2^{2/3 \cdot H(3/2 \cdot 1/226)} = o(1). \end{aligned}$$

Thus, a random A is good almost surely for a large enough n . We proceed with upper bounding $\text{bd}(G_{A, V-A})$ for a good A by counting the sets in \mathcal{N}_{V-A} , the family of neighbourhoods of subsets of $V-A$ in A .

Recall that $|V-A| \approx 2n/3$. The sets $S \subset V-A$ of size $i < n/226$ may contribute only as many as $\sum_{i=0}^{n/226} \binom{2n/3}{i}$ distinct neighbourhoods in A . The contribution of sets $S \subset V-A$ of size $\geq n/226$ may be bounded as follows. In each such S mark an arbitrary subset $X \subseteq S$ of size precisely $\lceil n/226 \rceil$. Call two large S 's equivalent if the same X was marked in both of them. Then, since every X sees at least αn vertices n , the contribution of the entire equivalence class of large sets defined X is at most $2^{n/3 - \alpha n}$. The number of X 's is at most $\binom{2n/3}{n/226}$. By plugging in the numerical value of α and using the entropy bound for $\binom{2n/3}{n/226}$, the entire contribution can be bounded by:

$$|\mathcal{N}_{V-A}| \leq \sum_{i=0}^{n/226} \binom{2n/3}{i} + \binom{2n/3}{n/226} \cdot 2^{n/3 - \alpha n} \leq 2^{0.3286n} \leq 2^{\frac{n}{3} - \frac{n}{226}}$$

the upper bound on $\text{bd}(G_{A, V-A}) = \log_2 |\mathcal{N}_{V-A}|$ follows. \square

As an immediate consequence of Lemma 3 and Proposition 3 we get

Theorem 7. *For any graph G , it holds that $\text{bw}(G) \leq \frac{n}{3} - \frac{n}{672} + o(n)$.*

6 Conclusion

Our results are the first non-trivial upper bounds on the value of a graph width parameter that hold *for every graph*. In this paper we gave three techniques to show such bounds, respectively Theorems 5, 6 and 7. At the moment the first two work only for linear boolean-width and the third is here applied only to boolean-width but it should work also for the linear case. We believe these bounds can be substantially improved. Combining Corollary 1 with Proposition 4 and Theorem 5 we can solve MAXIMUM WEIGHT INDEPENDENT SET and COUNTING INDEPENDENT SETS OF SIZE k in time $O^*(1.4108^n)$, and solve MINIMUM WEIGHT DOMINATING SET, MINIMUM WEIGHT TOTAL DOMINATING SET, MAXIMUM/MINIMUM WEIGHT INDEPENDENT DOMINATING SET, and counting versions of these, in time $O^*(1.9904^n)$. These runtimes beat brute-force search but faster algorithms exist in the literature, see [4]. Our goal was mainly to prove the viability of this new line of research by establishing structural qualitative results. The natural directions for further work are to improve the bounds and hence the runtime, and to increase the class of problems handled by Corollary 1.

References

1. Rémy Belmonte and Martin Vatshelle. Graph classes with structured neighborhoods and algorithmic applications. *TCS*, 2013. <http://dx.doi.org/10.1016/j.tcs.2013.01.011>.
2. Binh-Minh Bui-Xuan, Jan Arne Telle, and Martin Vatshelle. Boolean-width of graphs. *Theoretical Computer Science*, 412(39):5187–5204, 2011.
3. Vojtech Rödl Dwight Duffus, Peter Frankl. Maximal independent sets in bipartite graphs obtained from boolean lattices. *Eur. J. Comb.*, 32(1):1–9, 2011.
4. Fedor V. Fomin and Dieter Kratsch. *Exact Exponential Algorithms*. Texts in Theoretical Computer Science. 1st edition edition, 2010.
5. Zoltán Füredi. The number of maximal independent sets in connected graphs. *Journal of Graph Theory*, 11(4):463–470, 1987.
6. Petr Hliněný and Sang il Oum. Finding branch-decompositions and rank-decompositions. *SIAM J. Comput.*, 38(3):1012–1032, 2008.
7. Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, March 1963.
8. L. Ilinca and J. Kahn. Counting maximal antichains and independent sets. *Order*, 30(2):427–435, 2013.
9. Ki Hang Kim. *Boolean matrix theory and its applications*. Monographs and textbooks in pure and applied mathematics. Marcel Dekker, 1982.
10. Salil P. Vadhan. The complexity of counting in sparse, regular, and planar graphs. *SIAM Journal on Computing*, 31:398–427, 1997.
11. Martin Vatshelle. *New Width Parameters of Graphs*. PhD thesis, University of Bergen, 2012. ISBN:978-82-308-2098-8.

7 Appendix

Theorem 6 Let each vertex of V be assigned randomly and independently either to side A or to side B , resulting in $H(A, B, E'')$. Then it holds almost surely that $\text{co-im}(H) \geq \frac{1}{28}n - o(n)$. Consequently, $\text{co-bd}(H) \geq \frac{1}{801}n - o(n)$.

Proof. (Sketch) Consider the maximal packing \mathcal{P} of P_2 's and C_3 's in G . Let p be the number of P_2 's and c the number of C_3 's in \mathcal{P} . In H , about a quarter of P_2 's in \mathcal{P} will survive, and about half will become a single edge. About three quarters of C_3 's in \mathcal{P} will become (new) P_2 's. Let \mathcal{P}' be the packing of the surviving and the newly created P_2 's, and let \mathcal{R}' be the packing of edges resulting from the half-surviving P_2 's.

Let G' be the induced subgraph of G on $U = V - V(\mathcal{P})$. As before, G' consists of isolated vertices and edges, and about half of the latter will not survive into $H' = H|_U$.

Qualitatively, the argument is based on the fact that either \mathcal{P}' or $E(H')$ is large, each causing a loss in $\text{im}(G)$. What follows is an elaboration of this basic idea.

Since any induced matching M^* in H must miss at least one vertex for each P_2 in \mathcal{P}' , we conclude that

$$2 \text{co-im}(H) = n - 2 \text{im}(G) \geq |\mathcal{P}'| \geq \frac{1}{4}p + \frac{3}{4}c - o(n). \quad (5)$$

On the other hand, G' contains at least about $\frac{1}{2}|U|$ isolated vertices that can only be matched into $V(\mathcal{P})$. Call them U' . For every $P_2 \in \mathcal{P}'$, at least one of its vertices will not be matched to U' . The same applies to every $e \in \mathcal{R}'$. A closer look reveals that even if one of the endpoints of e is matched to U' , the other endpoint becomes unmatched. Therefore, the number of unmatched vertices in any M^* is at least about $|U'| - |V(\mathcal{P})| + |\mathcal{P}'| + |V(\mathcal{R}')|$. Thus,

$$\begin{aligned} 2 \text{co-im}(H) &\geq (n - 3(p + c))/2 - 3(p + c) + \left\{ \frac{1}{4}p + \frac{3}{4}c \right\} + 2 \cdot \frac{1}{2}p - o(n) = \\ &= n/2 - 3.25p - 3.75c - o(n). \end{aligned} \quad (6)$$

Optimizing the system of inequalities (5),(6), we conclude that $\text{co-im}(H) \geq \frac{1}{56}n - o(n)$, which is achieved at $p = \frac{1}{7}n$, $c = 0$. By Theorem 2, we conclude that $\text{co-bd}(H) \geq \frac{1}{56 \cdot 14.3}n - o(n) = \frac{1}{801}n - o(n)$. \square