

A width parameter useful for chordal and co-comparability graphs

Dong Yeap Kang^{*1}, O-joung Kwon^{†2}, Torstein J. F. Strømme³, and Jan Arne Telle³

¹Department of Mathematical Sciences, KAIST, Daejeon, South Korea.

²Logic and Semantics, Technische Universität Berlin, Berlin, Germany.

³Department of Informatics, University of Bergen, Norway.

February 11, 2017

Abstract

Belmonte and Vatshelle (TCS 2013) used mim-width, a graph width parameter bounded on interval graphs and permutation graphs, to explain existing algorithms for many domination-type problems on those graph classes. We investigate new graph classes of bounded mim-width, strictly extending interval graphs and permutation graphs. The graphs $K_t \boxplus K_t$ and $K_t \boxplus S_t$ are graphs obtained from the disjoint union of two cliques of size t , and one clique of size t and one independent set of size t respectively, by adding a perfect matching. We prove that :

- interval graphs are $(K_3 \boxplus S_3)$ -free chordal graphs; and $(K_t \boxplus S_t)$ -free chordal graphs have mim-width at most $t - 1$,
- permutation graphs are $(K_3 \boxplus K_3)$ -free co-comparability graphs; and $(K_t \boxplus K_t)$ -free co-comparability graphs have mim-width at most $t - 1$,
- chordal graphs and co-comparability graphs have unbounded mim-width in general.

We obtain several algorithmic consequences; for instance, while MINIMUM DOMINATING SET is NP-complete on chordal graphs, it can be solved in time $n^{\mathcal{O}(t)}$ on $(K_t \boxplus S_t)$ -free chordal graphs. The third statement strengthens a result of Belmonte and Vatshelle stating that either those classes do not have constant mim-width or a decomposition with constant mim-width cannot be computed in polynomial time unless $P = NP$.

We generalize these ideas to bigger graph classes. We introduce a new width parameter *sim-width*, of stronger modelling power than mim-width, by making a small change in the definition of mim-width. We prove that chordal graphs and co-comparability graphs have sim-width at most 1. We investigate a way to bound mim-width for graphs of bounded sim-width by excluding $K_t \boxplus K_t$ and $K_t \boxplus S_t$ as induced minors or induced subgraphs, and give algorithmic consequences. Lastly, we show that circle graphs have unbounded sim-width, and thus also unbounded mim-width.

*Supported by TJ Park Science Fellowship of POSCO TJ Park Foundation.

†Supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC consolidator grant DISTRUCT, agreement No. 648527).

⁰E-mail addresses: dynamical@kaist.ac.kr (D. Kang), ojoungkwon@gmail.com (O. Kwon), torstein.stromme@iib.uib.no (T. Strømme), Jan.Arne.Telle@uib.no (J. A. Telle).

An extended abstract appeared in the proceedings of 11th International Workshop on Algorithms and Computation, 2017.

1 Introduction

The study of structural graph “width” parameters like tree-width and clique-width have been ongoing since at least the 1990’s, and their algorithmic use has been steadily increasing [9]. A parallel and somewhat older field of study gives algorithms for special graph classes, such as chordal graphs and permutation graphs. Since the introduction of algorithms based on tree-width, which generalized algorithms for trees and series-parallel graphs, these two fields have been connected. Recently the fields became even more intertwined with the introduction of mim-width in 2012 [32], which yielded generalized algorithms for quite large graph classes. In the context of parameterized complexity a negative relationship holds between the modeling power of graph width parameters, i. e. what graph classes have bounded parameter values, and their analytical power, i. e. what problems become FPT or XP. For example, the family of graph classes of bounded width is strictly larger for clique-width than for tree-width, while under standard complexity assumptions the class of problems solvable in FPT time on a decomposition of bounded width is strictly larger for tree-width than for clique-width. For a parameter like mim-width its algorithmic use must therefore be carefully evaluated against its modelling power, which is much stronger than clique-width.

A common framework for defining graph width parameters is by branch decompositions over the vertex set. This is a natural hierarchical clustering of a graph G , represented as an unrooted binary tree T with the vertices of G at its leaves. Any edge of the tree defines a cut of G given by the leaves of the two subtrees that result from removing the edge from T . Judiciously choosing a cut-function to measure the complexity of such cuts, or rather of the bipartite subgraphs of G given by the edges crossing the cuts, this framework then defines a graph width parameter by a minmax relation, minimum over all trees and maximum over all its cuts. Restricting T to a path with added leaves we get a linear variant. The first graph parameter defined this way was carving-width [29], whose cut-function is simply the number of edges crossing the cut, and whose modelling power is weaker than tree-width. The carving-width of a graph G is thus the minimum, over all such branch decomposition trees, of the maximum number of edges crossing a cut given by an edge of the tree. Several graph width parameters have been defined this way. For example the mm-width [32], whose cut function is the size of the maximum matching and has modelling power equal to tree-width; and the rank-width [25], whose cut function is the GF[2]-rank of the adjacency matrix and has modelling power equal to clique-width.

This framework was used by Vatshelle in 2012 [32] to define the parameter mim-width whose cut function is the size of the maximum induced matching of the graph crossing the cut. Note that low mim-width allows quite complex cuts. Carving-width one allows just a single edge, mm-width one a star graph, and rank-width one a complete bipartite graph with some isolated vertices. In contrast, mim-width one allows any cut where the neighborhoods of the vertices in a color class can be ordered linearly w.r.t. inclusion. The modelling power of mim-width is much stronger than clique-width. Belmonte and Vatshelle showed that interval graphs and permutation graphs have linear mim-width at most one, and circular-arc graphs and trapezoid graphs have linear mim-width at most two [3]¹, whereas the clique-width of such graphs can be proportional to the square root of the number of vertices. With such strong modelling power it is clear that the analytical power of mim-width must suffer. The cuts in a decomposition of constant mim-width are too complex to allow FPT algorithms for interesting NP-hard problems. Instead, what we get is XP algorithms, for

¹In [3], all the related results are stated in terms of d -neighborhood equivalence, but in the proof, they actually gave a bound on mim-width or linear mim-width.



Figure 1: $K_5 \boxplus S_5$ and $K_5 \boxplus K_5$.

the class of LC-VSVP problems [8] - locally checkable vertex subset and vertex partitioning problems - defined in Section 2.4. For classes of bounded mim-width this gives a common explanation for many classical results in the field of algorithms on special graph classes.

In this paper we extend these results on mim-width in several ways. Firstly, we show that chordal graphs and co-comparability graphs have unbounded mim-width, thereby answering a question of [3]², and giving evidence to the intuition that mim-width is useful for large graph classes having a linear structure rather than those having a tree-like structure. Secondly, by excluding certain subgraphs, like $K_t \boxplus S_t$ obtained from the disjoint union of a clique of size t and an independent set of size t by adding a perfect matching, we find subclasses of chordal graphs and co-comparability graphs for which we can nevertheless compute a bounded mim-width decomposition in polynomial time and thereby solve LC-VSVP problems. See Figure 1 for illustrations. Note that already for $K_3 \boxplus S_3$ -free chordal graphs the tree-like structure allowed by mim-width is necessary, as they have unbounded linear mim-width. Thirdly, we introduce sim-width, of modeling power even stronger than mim-width, and start a study of its properties.

The graph width parameter sim-width is defined within the same framework as mim-width with only a slight change to its cut function, simply requiring that a special induced matching across a cut cannot contain edges between any pair of vertices on the same side of the cut. This exploits the fact that a cut function for branch decompositions over the vertex set of a graph need not be a parameter of the bipartite graph on edges crossing the cut. The cuts allowed by sim-width are more complex than for mim-width, making sim-width applicable to well-known graph classes having a tree-like structure. We show that chordal graphs and co-comparability graphs have sim-width one and that these decompositions can be found in polynomial time. See Figure 2 for an inclusion diagram of some well-known graph classes illustrating these results. Since LC-VSVP problems like MINIMUM DOMINATING SET are NP-complete on chordal graphs we cannot expect XP algorithms parameterized by sim-width. However, for graphs of sim-width w , when excluding subgraphs $K_t \boxplus K_t$ or $K_t \boxplus S_t$, either as induced subgraphs or induced minors, we get graphs of bounded mim-width. The induced minor relation is natural since graphs of bounded sim-width are closed under induced minors, which might be of independent interest when taking the structural point of view. We show that by the alternate parametrization $w + t$ the LC-VSVP problems are solvable in XP time on graphs excluding $K_t \boxplus K_t$ or $K_t \boxplus S_t$, when given with a decomposition of sim-width w . The class of circle graphs is one of the classes listed in [3] where either graphs in the class do not have constant mim-width, or it is NP-complete to find such a decomposition. Using a technique recently introduced by Mengel [24] we prove that sim-width of circle graphs is unbounded, which implies that mim-width of circle graphs is also unbounded.

Let us mention some more related work. Golovach et al [18] applied linear mim-width in the context of enumeration algorithms, showing that all minimal dominating sets of graphs of bounded

²Our result appeared on arXiv June 2016. In August 2016 a similar result by Mengel, developed independently, also appeared on arXiv [24].




Figure 2: Inclusion diagram of some well-known graph classes. (I) Classes where clique-width and rank-width are constant. (II) Classes where mim-width is constant. (III) Classes where sim-width is constant. (IV) Classes where it is unknown if sim-width is constant. (V) Classes where sim-width is unbounded.

linear mim-width can be enumerated with polynomial delay using polynomial space. The graph parameter mim-width has also been used for knowledge compilation and model counting in the field of satisfiability [7, 27, 15]. The LC-VSVP problems include the class of domination-type problems known as (σ, ρ) -DOMINATION problems, whose intractability on chordal graphs is well known [6]. For two subsets of natural numbers σ, ρ a set S of vertices is called (σ, ρ) -dominating if for every vertex $v \in S$, $|S \cap N(v)| \in \sigma$, and for every $v \notin S$, $|S \cap N(v)| \in \rho$. Golovach et al [17] showed that for fixed σ, ρ the problem of deciding if a given chordal graph has a (σ, ρ) -dominating set, is NP-complete whenever there exists at least one chordal graph containing more than one (σ, ρ) -dominating set, as this graph can then be used as a gadget in a reduction. Golovach, Kratochvíl and Suchý [19] extended these results to the parameterized setting, showing that the existence of a (σ, ρ) -dominating set of size k , and at most k , are W[1]-complete problems when parameterized by k for any pair of finite sets σ and ρ . In contrast, combining our bounds on mim-width and algorithms of Bui-Xuan, Telle, and Vatshelle [8] we obtain the following.

Theorem 1.1. *Let $t \geq 2$ be an integer. Given an n -vertex $(K_t \boxminus S_t)$ -free chordal graph or an n -vertex $(K_t \boxminus K_t)$ -free co-comparability graph, every fixed LC-VSVP problem can be solved in time $n^{O(t)}$.*

As a specific example, we show that MINIMUM DOMINATING SET can be solved in time $O(n^{3t+4})$ and q -COLORING can be solved in time $O(qn^{3qt+4})$. We note that given an n -vertex graph, one can test in time $O(n^{2t})$ whether it contains an induced subgraph isomorphic to $K_t \boxminus S_t$ or not. Therefore, a membership testing for $(K_t \boxminus S_t)$ -free chordal graphs and the algorithm in Theorem 1.1

can be applied in time $n^{\mathcal{O}(t)}$. The same argument holds for $(K_t \boxminus K_t)$ -free co-comparability graphs.

The remainder of this paper is organized as follows. Section 2 contains all the necessary notions required for our results. In Section 3, we prove that chordal graphs have sim-width at most 1 and mim-width at most $t - 1$ if they are $(K_t \boxminus S_t)$ -free. Similarly we show that co-comparability graphs have linear sim-width at most 1 and linear mim-width at most $t - 1$ if they are $(K_t \boxminus K_t)$ -free. We provide polynomial-time algorithms to find such decompositions, and discuss their algorithmic consequences for LC-VSVP problems. We also show that chordal graphs and co-comparability graphs have unbounded mim-width. In Section 4, we bound the mim-width of graphs of sim-width w that do not contain $K_t \boxminus K_t$ and $K_t \boxminus S_t$ as induced subgraphs or induced minors, and give algorithmic consequences. We also show that graphs of bounded sim-width are closed under induced minors. Lastly, we show in Section 5 that circle graphs have unbounded sim-width. We finish with some research questions related to sim-width in Section 6.

2 Preliminaries

We denote the vertex set and edge set of a graph G by $V(G)$ and $E(G)$, respectively. Let G be a graph. For a vertex v of G , we denote by $N_G(v)$ the set of neighbors of v in G . For $v \in V(G)$ and $X \subseteq V(G)$, we denote by $G - v$ the graph obtained from G by removing v , and denote by $G - X$ the graph obtained from G by removing all vertices in X . For $e \in E(G)$, we denote by $G - e$ the graph obtained from G by removing e , and denote by G/e the graph obtained from G by contracting e . For a vertex v of G with exactly two neighbors v_1 and v_2 that are non-adjacent, the operation of removing v and adding the edge v_1v_2 is called *smoothing* the vertex v . For $X \subseteq V(G)$, we denote by $G[X]$ the subgraph of G induced by X . A *clique* in G is a set of vertices of G that are pairwise adjacent, and an *independent set* in G is a set of vertices that are pairwise non-adjacent. A set of edges $\{v_1w_1, v_2w_2, \dots, v_mw_m\}$ of G is called an *induced matching* in G if there are no other edges in $G[\{v_1, \dots, v_m, w_1, \dots, w_m\}]$. A matrix M is called the *adjacency matrix* of G if the rows and columns of M are indexed by $V(G)$, and for $v, w \in V(G)$, $M[v, w] = 1$ if v and w are adjacent in G , and $M[v, w] = 0$ otherwise.

A pair of vertex subsets (A, B) of a graph G is called a *vertex bipartition* if $A \cap B = \emptyset$ and $A \cup B = V(G)$. For a vertex bipartition (A, B) of a graph G , we denote by $G[A, B]$ the bipartite graph on the bipartition (A, B) where for $a \in A$ and $b \in B$, a and b are adjacent in $G[A, B]$ if and only if they are adjacent in G . For a vertex bipartition (A, B) of G and an induced matching $\{v_1w_1, v_2w_2, \dots, v_mw_m\}$ in G where $v_1, \dots, v_m \in A$ and $w_1, \dots, w_m \in B$, we say that it is an induced matching in G between A and B .

We denote by \mathbb{N} the set of all non-negative integers, and let $\mathbb{N}^+ := \mathbb{N} \setminus \{0\}$.

2.1 Graph classes

A tree is called *subcubic* if every internal node has exactly 3 neighbors. A tree T is called a *caterpillar* if it contains a path P such that every vertex in $V(T) \setminus V(P)$ has a neighbor in P . The complete graph on n vertices is denoted by K_n .

A graph is *chordal* if it contains no induced subgraph isomorphic to a cycle of length 4 or more. A graph is a *split* graph if it can be partitioned into two vertex sets C and I where C is a clique and I is an independent set. A graph is an *interval graph* if it is the intersection graph of a family of intervals on the real line. Every split graph and interval graph is chordal. An ordering v_1, \dots, v_n

of the vertex set of a graph G is called a *co-comparability ordering* if for every integers i, j, k with $1 \leq i < j < k \leq n$,

- if v_i is adjacent to v_k , then v_j is adjacent to v_i or v_k .

This condition is equivalent to the following: for every integers i, j, k with $1 \leq i < j < k \leq n$, v_j has a neighbor in every path from v_i to v_k avoiding v_j . A graph is a *co-comparability graph* if it admits a co-comparability ordering. Every co-comparability graph is the complement of some comparability graph, where comparability graphs are graphs that can be obtained from some partial order by connecting pairs of elements that are comparable to each other. A graph is a *permutation graph* if it is the intersection graph of line segments whose endpoints lie on two parallel lines. Permutation graphs are co-comparability graphs [11]. A graph is a *circle graph* if it is the intersection graph of chords in a circle.

For positive integer n , let $K_n \boxminus K_n$ be the graph on $\{v_1^1, \dots, v_n^1, v_1^2, \dots, v_n^2\}$ such that for all $i, j \in \{1, \dots, n\}$,

- $\{v_1^1, \dots, v_n^1\}$ and $\{v_1^2, \dots, v_n^2\}$ are cliques,
- v_i^1 is adjacent to v_j^2 if and only if $i = j$,

and let $K_n \boxminus S_n$ be the graph on $\{v_1^1, \dots, v_n^1, v_1^2, \dots, v_n^2\}$ such that for all $i, j \in \{1, \dots, n\}$,

- $\{v_1^1, \dots, v_n^1\}$ is a clique, $\{v_1^2, \dots, v_n^2\}$ is an independent set,
- v_i^1 is adjacent to v_j^2 if and only if $i = j$.

Since $K_2 \boxminus K_2$ is an induced cycle of length 4, chordal graphs do not contain $K_2 \boxminus K_2$ as an induced subgraph. We observe that $K_3 \boxminus S_3$ is not a co-comparability graph.

Lemma 2.1. *The graph $K_3 \boxminus S_3$ is not a co-comparability graph.*

Proof. Let G be a graph on $\{v_1, v_2, v_3\} \cup \{w_1, w_2, w_3\}$ such that

- $\{v_1, v_2, v_3\}$ is a clique, $\{w_1, w_2, w_3\}$ is an independent set, and
- v_i is adjacent to w_j if and only if $i = j$.

It is clear that G is isomorphic to $K_3 \boxminus S_3$. Suppose G admits a co-comparability ordering. By relabeling if necessary, we may assume w_1, w_2, w_3 appear in the co-comparability ordering in that order. However, there is a path $w_1 v_1 v_3 w_3$ from w_1 to w_3 avoiding v_2 and w_2 , and thus, it contradicts to the assumption. We conclude that $K_3 \boxminus S_3$ is not a co-comparability graph. \square

2.2 Graph relations

Let G be a graph. A graph H is a *subgraph* of G if H can be obtained from G by removing some vertices and edges. A graph H is an *induced subgraph* of G if $H = G[X]$ for some $X \subseteq V(G)$. A graph H is an *induced minor* of G if H can be obtained from G by a sequence of removing vertices and contracting edges. A graph H is a *minor* of G if H can be obtained from G by a sequence of removing vertices, removing edges, and contracting edges. For a graph H , we say a graph is *H -free* if it contains no induced subgraph isomorphic to H .

A *minor model* of a graph H in G is a function η with the domain $V(H) \cup E(H)$, where

- for every $v \in V(H)$, $\eta(v)$ is a non-empty connected subgraph of G , all pairwise vertex-disjoint
- for every edge e of H , $\eta(e)$ is an edge of G , all distinct
- if $e \in E(H)$ and $v \in V(H)$ then $\eta(e) \not\subseteq E(\eta(v))$,
- for every edge $e = uv$ of H , $\eta(e)$ has one end in $V(\eta(u))$ and the other in $V(\eta(v))$.

It is well known that H is a minor of G if and only if there is a minor model of H in G . A minor model η of a graph H in G is an *induced minor model* if for every distinct vertices v_1 and v_2 in H that are non-adjacent, there are no edges between $V(\eta(v_1))$ and $V(\eta(v_2))$. A graph H is an induced minor of G if and only if there is an induced minor model of H in G .

2.3 Width parameters

For sets A and B , a function $f : 2^A \rightarrow B$ is *symmetric* if for every $Z \subseteq A$, $f(Z) = f(A \setminus Z)$.

For a graph G and a vertex set $A \subseteq V(G)$, we define functions cutrk_G , mim_G , and sim_G from $2^{V(G)}$ to \mathbb{N} such that

- $\text{cutrk}_G(A)$ is the rank of the matrix $M[A, V(G) \setminus A]$ where M is the adjacency matrix of G and the rank is computed over the binary field,
- $\text{mim}_G(A)$ is the maximum size of an induced matching of $G[A, V(G) \setminus A]$,
- $\text{sim}_G(A)$ is the maximum size of an induced matching between A and $V(G) \setminus A$ in G .

For a graph G , a pair (T, L) of a subcubic tree T and a bijection L from $V(G)$ to the set of leaves of T is called a *branch-decomposition*. For each edge e of T , let T_1^e and T_2^e be the two connected components of $T - e$, and let (A_1^e, A_2^e) be the vertex bipartition of G such that for each $i \in \{1, 2\}$, A_i^e is the set of all vertices in G mapped to leaves contained in T_i^e by L . We call (A_1^e, A_2^e) the vertex bipartition of G associated with e . For a branch-decomposition (T, L) of a graph G and an edge e in T and a symmetric function $f : 2^{V(G)} \rightarrow \mathbb{N}$, the *f-width* of e is define as $f(A_1^e)$ where (A_1^e, A_2^e) is the vertex bipartition associated with e . The *f-width* of (T, L) is the maximum *f-width* over all edges in T , and the *f-width* of G is the minimum *f-width* over all branch-decompositions of G . If $|V(G)| \leq 1$, then G does not admit a branch-decomposition, and the *f-width* of G is defined to be 0.

The *rank-width* of a graph G , denoted by $\text{rw}(G)$, is the cutrk_G -width of G , and the *mim-width* of a graph G , denoted by $\text{mimw}(G)$, is the mim_G -width of G , and the *sim-width* of a graph G , denoted by $\text{simw}(G)$, is the sim_G -width of G . For convenience, the *f-width* of a branch-decomposition is also called a rank-width, mim-width, or sim-width depending on the function f .

If T is a subcubic caterpillar tree, then a branch-decomposition (T, L) is called a *linear branch-decomposition*. The *linear f-width* of G is the minimum *f-width* over all linear branch-decompositions of G . The *linear mim-width* of a graph G , denoted by $\text{lmimw}(G)$, is the linear mim_G -width of G , and the *linear sim-width* of a graph G , denoted by $\text{lsimw}(G)$, is the linear sim_G -width of G .

By definitions we have the following.

Lemma 2.2. *For a graph G , we have $\text{simw}(G) \leq \text{mimw}(G) \leq \text{rw}(G)$.*

We frequently use the following fact.

Lemma 2.3. *Let G be a graph, let w be a positive integer, and let $f : 2^{V(G)} \rightarrow \mathbb{N}$ be a symmetric function. If G has f -width at most w , then G admits a vertex bipartition (A_1, A_2) where $f(A_1) \leq w$ and for each $i \in \{1, 2\}$, $\frac{|V(G)|}{3} < |A_i| \leq \frac{2|V(G)|}{3}$.*

Proof. Let (T, L) be a branch-decomposition of G of f -width at most w . We subdivide an edge of T , and regard the new vertex as a root node. For each node $t \in V(T)$, let $\mu(t)$ be the number of leaves of T that are descendants of t . Now, we choose a node t that is farthest from the root node such that $\mu(t) > \frac{|V(G)|}{3}$. By the choice of t , for each child t' of t , $\mu(t') \leq \frac{|V(G)|}{3}$. Therefore, $\frac{|V(G)|}{3} < \mu(t) \leq \frac{2|V(G)|}{3}$. Let e be the edge connecting the node t and its parent. By the construction, the vertex bipartition associated with e is a required bipartition. \square

We also use tree-decompositions in Section 3. A *tree-decomposition* of a graph G is a pair $(T, \mathcal{B} = \{B_t\}_{t \in V(T)})$ such that

- (1) $\bigcup_{t \in V(T)} B_t = V(G)$,
- (2) for every edge in G , there exists B_t containing both end vertices,
- (3) for $t_1, t_2, t_3 \in V(T)$, $B_{t_1} \cap B_{t_3} \subseteq B_{t_2}$ whenever t_2 is on the path from t_1 to t_3 .

Each vertex subset B_t is called a *bag* of the tree-decomposition. The *width* of a tree-decomposition is $w - 1$ where w is the maximum size of a bag in the tree-decomposition, and the *tree-width* of a graph is the minimum width over all tree-decompositions of the graph. It is well known that a graph has tree-width at most w if and only if it is a subgraph of a chordal graph with maximum clique size at most $w + 1$; see for instance [5]. Furthermore, chordal graphs admit a tree-decomposition where each bag induces a maximal clique of the graph. We will use this fact in Section 3.

2.4 LC-VSVP problems

Telle and Proskurowski [31] classified a class of problems called *locally checkable vertex subset and vertex partitioning problems*, which is a subclass of MSO_1 problems. These problems generalize problems like MAXIMUM INDEPENDENT SET, MINIMUM DOMINATING SET, q -COLORING etc.

Let σ and ρ be finite or co-finite subsets of \mathbb{N} . For a graph G and $S \subseteq V(G)$, we call S a (σ, ρ) -*dominating set* of G if

- for every $v \in S$, $|N_G(v) \cap S| \in \sigma$, and
- for every $v \in V(G) \setminus S$, $|N_G(v) \cap S| \in \rho$.

For instance, a $(0, \mathbb{N})$ -set is an independent set as there are no edges inside of the set, and we do not care about the adjacency between S and $V(G) \setminus S$. Another example is that an $(\mathbb{N}, \mathbb{N}^+)$ -set is a dominating set as we require that for each vertex in $V(G) \setminus S$, it has at least one neighbor in S . See [31, Table 4.1] for more examples. The MIN-(OR MAX-) (σ, ρ) -DOMINATION problem is a problem to find a minimum (or maximum) (σ, ρ) -dominating set in an input graph G , and possibly on vertex-weighted graphs. These problems also called as *locally checkable vertex subset* problems.

For a positive integer q , a $(q \times q)$ -matrix D_q is called a *degree constraint* matrix if each element is either a finite or co-finite subset of \mathbb{N} . A partition $\{V_1, V_2, \dots, V_q\}$ of the vertex set of a graph G is called a D_q -*partition* if

- for every $i, j \in \{1, \dots, q\}$ and $v \in V_i$, $|N_G(v) \cap V_j| \in D_q[i, j]$.

For instance, if we take a matrix D_q where all diagonal entries are 0, and all other entries are \mathbb{N} , then a D_q -partition is a partition into q independent sets, which corresponds to a q -coloring of the graph. The D_q -PARTITIONING problem is a problem deciding if an input graph admits a D_q -partition or not. These problems are also called as *locally checkable vertex partitioning* problems.

All these problems will be called *locally checkable vertex subset and vertex partitioning problems*, shortly LC-VSVP problems. As shown in [8] the runtime solving an LC-VSVP problem by dynamic programming relates to the finite or co-finite subsets of \mathbb{N} used in its definition. The following function d is central.

1. Let $d(\mathbb{N}) = 0$.
2. For every finite or co-finite set $\mu \subseteq \mathbb{N}$, let $d(\mu) = 1 + \min(\max\{x \in \mathbb{N} : x \in \mu\}, \max\{x \in \mathbb{N} : x \notin \mu\})$.

For example, for MINIMUM DOMINATING SET and q -COLORING problems we plug in $d = 1$ because $\max(d(\mathbb{N}), d(\mathbb{N}^+)) = 1$ and $\max(d(0), d(\mathbb{N})) = 1$.

Belmonte and Vatshelle [3] proved the following application of mim-width.

Theorem 2.4 (Belmonte and Vatshelle [3] and Bui-Xuan, Telle, and Vatshelle [8]). *(1) Given an n -vertex graph and its branch-decomposition (T, L) of mim-width w and $\sigma, \rho \subseteq \mathbb{N}$, one can solve MIN-(OR MAX-)(σ, ρ)-DOMINATION in time $\mathcal{O}(n^{3dw+4})$ where $d = \max(d(\sigma), d(\rho))$.*

(2) Given an n -vertex graph and its branch-decomposition (T, L) of mim-width w and a $(q \times q)$ -matrix D_q , one can solve D_q -PARTITIONING in time $\mathcal{O}(qn^{3dwq+4})$ where $d = \max_{i,j} d(D_q[i, j])$.

3 Mim-width of chordal graphs and co-comparability graphs

In this section, we show that chordal graphs admit a branch-decomposition (T, L) such that

1. (T, L) has sim-width at most 1, and
2. (T, L) has mim-width at most $t - 1$ if the given graph is $(K_t \boxminus S_t)$ -free for some $t \geq 2$.

We combine the second statement with Theorem 2.4, and show that LC-VSVP problems can be efficiently solved for $(K_t \boxminus S_t)$ -free chordal graphs (Corollary 3.2). In the same context, we show that co-comparability graphs admit a linear branch-decomposition such that

1. (T, L) has linear sim-width at most 1, and
2. (T, L) has linear mim-width at most $t - 1$ if the given graph is $(K_t \boxminus K_t)$ -free for some $t \geq 2$.

We combine the second statement with Theorem 2.4, and show that LC-VSVP problems can be efficiently solved for $(K_t \boxminus K_t)$ -free co-comparability graphs (Corollary 3.7). We also prove that chordal graphs and co-comparability graphs have unbounded mim-width in general.




Figure 3: Constructing a branch-decomposition (T, L) of a chordal graph G of sim-width at most 1 from its tree-decomposition.

3.1 Mim-width of chordal graphs

We prove the following.

Proposition 3.1. *Given a chordal graph G , one can in time $\mathcal{O}(|V(G)| + |E(G)|)$ output a branch-decomposition (T, L) with the following property:*

- (T, L) has sim-width at most 1,
- (T, L) has mim-width at most $t - 1$ if G is $(K_t \boxminus S_t)$ -free for some integer $t \geq 2$.

To show Proposition 3.1, we use the fact that chordal graphs admit a tree-decomposition whose bags are maximal cliques. Such a tree-decomposition can be easily transformed from a *clique-tree* of a chordal graph. A *clique-tree* of a chordal graph G is a pair $(T, \{C_t\}_{t \in V(T)})$, where

- (1) T is a tree,
- (2) each C_t is a maximal clique of G , and
- (3) for each $v \in V(G)$, the vertex set $\{t \in V(T) : v \in C_t\}$ induces a subtree of T .

Gavril [16] showed that chordal graphs admit a clique-tree, and it is known that given a chordal graph G , its clique-tree can be constructed in time $\mathcal{O}(|V(G)| + |E(G)|)$ [4, 21].

Proof of Proposition 3.1. Let G be a chordal graph. We compute a tree-decomposition $(F, \mathcal{B} = \{B_t\}_{t \in V(F)})$ of G whose bags induce maximal cliques of G in time $\mathcal{O}(|V(G)| + |E(G)|)$. Let us choose a root node r of F .

We construct a tree (T, L) from F as follows.

1. We attach a leaf r' to the root node r of F and regard it as the parent of r and let $B_{r'} := \emptyset$.
2. For every $t \in V(F)$ with its parent t' , we subdivide the edge tt' into a path $tv_1^t \cdots v_{|B_t \setminus B_{t'}|}^t t'$, and for each $j \in \{1, \dots, |B_t \setminus B_{t'}|\}$, we attach a leaf z_j^t to v_j^t and assign the vertices of $B_t \setminus B_{t'}$ to $L(z_1^t), \dots, L(z_{|B_t \setminus B_{t'}|}^t)$ in any order. Then remove r' .
3. We transform the resulting tree into a tree of maximum degree at most 3. For every $t \in V(F)$, we do the following. Let t_1, \dots, t_m be the children of t in F . We remove t and introduce a path $w_1^t w_2^t \cdots w_m^t$. If t is a leaf, then we just remove it. We add an edge $w_1^t v_1^t$, and for each $i \in \{1, \dots, m\}$, add an edge $w_i^t v_{|B_{t_i} \setminus B_t}^{t_i}$.

4. Let T' be the resulting tree, and we obtain a tree T from T' by smoothing all nodes of degree 2. Let (T, L) be the resulting branch-decomposition. See Figure 3 for an illustration of the construction.

We can construct (T, L) in linear time as the number of nodes in T is $\mathcal{O}(|V(G)|)$. We consider T as a rooted tree with the root $z_{B_r \setminus B_{r'}}^r$.

We claim that (T, L) has sim-width at most 1. We prove a stronger result that for every edge e of T with a vertex bipartition (A, B) associated with e , either $N_G(A) \cap B$ or $N_G(B) \cap A$ is a clique.

Claim 1. *Let e be an edge of T and let (A, B) be the vertex bipartition of G associated with e . Then either $N_G(A) \cap B$ or $N_G(B) \cap A$ is a clique.*

Proof. For convenience, we prove for T' , which is the tree before smoothing. We may assume that both end nodes of e are internal nodes of T' , otherwise, it is trivial. There are four types of e :

1. $e = v_i^t v_{i+1}^t$ for some $t \in V(F)$, its parent t' , and $i \in \{1, \dots, |B_t \setminus B_{t'}| - 1\}$.
2. $e = w_1^t v_1^t$ for some $t \in V(F)$.
3. $e = v_{|B_{t_i} \setminus B_t|}^{t_i} w_i^t$ for some $t \in V(F)$ and its child t_i .
4. $e = w_i^t w_{i+1}^t$ for some $t \in V(F)$ and its child t_i .

Suppose $e = v_i^t v_{i+1}^t$ for some t and i , and let t' be the parent of t . We may assume that A is the set of all vertices assigned to the descendants of v_i^t . Note that $B_t \cap B_{t'}$ separates A and $B \setminus B_t$ in G . Therefore, for each $v \in A$, $N_G(v) \cap B$ is contained in B_t , and $N_G(A) \cap B$ is a clique. We can similarly prove for Cases 2 and 3.

Now, let t_1, \dots, t_m be the children of t , and let w_1^t, \dots, w_m^t be the vertices that were replaced from t in the algorithm. We assume $e = w_i^t w_{i+1}^t$ for some $i \in \{1, \dots, m-1\}$. For each $j \in \{1, \dots, m\}$, let L_j be the set of all vertices assigned to the descendants of $v_{|B_{t_j} \setminus B_t|}^{t_j}$. Without loss of generality, we may assume that $B_t \subseteq B$. We can observe that $A = \bigcup_{i+1 \leq j \leq m} L_j$.

Let j_1 and j_2 be integers such that $1 \leq j_1 \leq i < j_2 \leq m$. Note that B_t separates L_{j_1} and L_{j_2} in G . Therefore, for every $v \in A$, $N_G(v) \cap B$ is contained in B_t , and $N_G(A) \cap B$ is a clique, as required. \diamond

We prove that when G is $(K_t \boxplus S_t)$ -free for some $t \geq 2$, (T, L) has mim-width at most $t-1$.

Claim 2. *Let $t \geq 2$ be an integer. If G is $(K_t \boxplus S_t)$ -free, then (T, L) has mim-width at most $t-1$.*

Proof. We show that (T, L) has mim-width at most $t-1$. Suppose for contradiction that there is an edge e of T with the vertex bipartition (A, B) of G associated with e such that $\text{mim}_G(A) \geq t$. We may assume both end nodes of e are internal nodes of T .

By Claim 1, one of $N_G(A) \cap B$ and $N_G(B) \cap A$ is a clique. Without loss of generality we assume $N_G(B) \cap A$ is a clique C . Since $\text{mim}_G(A) \geq t$, there is an induced matching $\{a_1 b_1, \dots, a_t b_t\}$ in $G[A, B]$ where $a_1, \dots, a_t \in A$. Since $N_G(B) \cap A$ is a clique C , there are no edges between vertices in $\{b_1, \dots, b_t\}$, otherwise, it creates an induced cycle of length 4. Thus, we have an induced subgraph isomorphic to $K_t \boxplus S_t$, which contradicts to our assumption. We conclude that (T, L) has mim-width at most $t-1$. \diamond

This concludes the proposition. \square

As a corollary of Proposition 3.1, we obtain the following.

Corollary 3.2. *Let $t \geq 2$ be an integer.*

- (1) *Given an n -vertex $(K_t \boxminus S_t)$ -free chordal graph and $\sigma, \rho \subseteq \mathbb{N}$, one can solve MIN-(OR MAX-)
PARTITIONING in time $\mathcal{O}(n^{3d(t-1)+4})$ where $d = \max(d(\sigma), d(\rho))$.*
- (2) *Given an n -vertex $(K_t \boxminus S_t)$ -free chordal graph and a $(q \times q)$ -matrix D_q , one can solve D_q -
PARTITIONING in time $\mathcal{O}(qn^{3dq(t-1)+4})$ where $d = \max_{i,j} d(D_q[i, j])$.*

This generalizes the algorithms for interval graphs, as interval graphs are $(K_3 \boxminus S_3)$ -free chordal graphs [22].

We now prove the lower bound on the mim-width of general chordal graphs. We show this for split graphs, which form a subclass of the class of chordal graphs. The Sauer-Shelah lemma [28, 30] is essential in the proof.

Theorem 3.3 (Sauer-Shelah lemma [28, 30]). *Let t be a positive integer and let M be an $X \times Y$ $(0, 1)$ -matrix such that $|Y| \geq 2$ and any two row vectors of M are distinct. If $|X| \geq |Y|^t$, then there are $X' \subseteq X$, $Y' \subseteq Y$ such that $|X'| = 2^t$, $|Y'| = t$, and all possible row vectors of length t appear in $M[X', Y']$.*

Proposition 3.4. *For every large enough n , there is a split graph on n vertices having mim-width at least $\frac{\log_2 n}{5 \log_2(\log_2 n)}$.*

Proof. Let $m \geq 4$ be an integer and let $n := m + (2^m - 1)$. Let G be a split graph on the vertex bipartition (C, I) where C is a clique of size m , I is an independent set of size $2^m - 1$, and all vertices in I have pairwise distinct non-empty neighborhoods on C . We claim that every branch-decomposition of G has mim-width at least $\frac{\log_2 n}{5 \log_2(\log_2 n)}$.

Let (T, L) be a branch-decomposition of G . By Lemma 2.3, there is a vertex bipartition (A_1, A_2) of G associated with e satisfies that for each $i \in \{1, 2\}$, $\frac{n}{3} < |A_i| \leq \frac{2n}{3}$. Without loss of generality, we may assume that $|A_1 \cap C| \geq |A_2 \cap C|$, and thus we have $\frac{m}{2} \leq |A_1 \cap C| \leq m$ and $|A_2 \cap C| \leq \frac{m}{2}$.

Claim 3. *There are at least $|A_1 \cap C|^{\frac{m}{4 \log_2 m}}$ vertices in $A_2 \cap I$ that have pairwise distinct neighbors in $A_1 \cap C$.*

Proof. Since $m \geq 4$, we have $|A_2 \cap I| = |A_2| - |A_2 \cap C| > \frac{n}{3} - \frac{m}{2} \geq \frac{2^m - \frac{m}{2} - 1}{3} \geq \frac{1}{4} 2^m$. Since $|A_2 \cap C| < \frac{m}{2}$, there are at least

$$\frac{1}{4} \cdot \frac{2^m}{2^{|A_2 \cap C|}} \geq \frac{1}{4} 2^{\frac{m}{2}}$$

vertices in $A_2 \cap I$ that have pairwise distinct neighbors on $A_1 \cap C$. Thus, we have

$$\frac{1}{4} 2^{\frac{m}{2}} \geq m^{\frac{m}{4 \log_2 m}} \geq |A_1 \cap C|^{\frac{m}{4 \log_2 m}},$$

as required. \diamond

Let $I' \subseteq A_2 \cap I$ be a maximal set of vertices that have pairwise distinct neighbors in $A_1 \cap C$. By the Sauer-Shelah lemma, there is an induced matching of size $\frac{m}{4 \log_2 m}$ between $A_1 \cap C$ and I' . It implies that (T, L) has mim-width at least $\frac{m}{4 \log_2 m}$. As (T, L) was chosen arbitrary, the mim-width of G is at least $\frac{m}{4 \log_2 m} \geq \frac{(\log_2 n) - 1}{4 \log_2(\log_2 n)} \geq \frac{\log_2 n}{5 \log_2(\log_2 n)}$. \square

3.2 Mim-width of co-comparability graphs

We observe similar properties for co-comparability graphs.

Theorem 3.5 (McConnell and Spinrad [23]). *Given a co-comparability graph G , one can output a co-comparability ordering in time $\mathcal{O}(|V(G)| + |E(G)|)$.*

Proposition 3.6. *Given a co-comparability graph G , one can in time $\mathcal{O}(|V(G)| + |E(G)|)$ output a linear branch-decomposition (T, L) with the following property:*

- (T, L) has linear sim-width at most 1,
- (T, L) has linear mim-width at most $t - 1$ if G is $(K_t \boxminus K_t)$ -free for some integer $t \geq 2$.

Proof. Let G be a co-comparability graph. Using Theorem 3.5, we can obtain its co-comparability ordering v_1, \dots, v_n in time $\mathcal{O}(|V(G)| + |E(G)|)$. From this, we take a linear branch-decomposition (T, L) following the co-comparability ordering. We claim that for each $i \in \{2, \dots, n-1\}$, there is no induced matching of size 2 between $\{v_1, \dots, v_i\}$ and $\{v_{i+1}, \dots, v_n\}$. Suppose there are $i_1, i_2 \in \{1, \dots, i\}$ and $j_1, j_2 \in \{i+1, \dots, n\}$ such that $\{v_{i_1}v_{j_1}, v_{i_2}v_{j_2}\}$ is an induced matching of G . Without loss of generality we may assume that $i_1 < i_2$. Then we have $i_1 < i_2 < j_1$, and thus by the definition of the co-comparability ordering, v_{i_2} should be adjacent to one of v_{i_1} and v_{j_1} , which contradicts to our assumption. Therefore, there is no induced matching of size 2 between $\{v_1, \dots, v_i\}$ and $\{v_{i+1}, \dots, v_n\}$. It implies that (T, L) has sim-width at most 1.

Now, suppose that G is $(K_t \boxminus K_t)$ -free for $t \geq 2$. We prove that for every $i \in \{1, \dots, n\}$, there are no induced matchings of size t in $G[\{v_1, \dots, v_i\}, \{v_{i+1}, \dots, v_n\}]$. For contradiction, suppose there is an induced matching $\{v_{i_1}v_{j_1}, \dots, v_{i_t}v_{j_t}\}$ in $G[\{v_1, \dots, v_i\}, \{v_{i+1}, \dots, v_n\}]$ where $i_1, \dots, i_t \in \{1, \dots, i\}$. We claim that $\{v_{i_1}, \dots, v_{i_t}\}$ and $\{v_{j_1}, \dots, v_{j_t}\}$ are cliques.

Claim 4. $\{v_{i_1}, \dots, v_{i_t}\}$ and $\{v_{j_1}, \dots, v_{j_t}\}$ are cliques.

Proof. Let $x, y \in \{1, \dots, t\}$. Assume that $i_x < i_y$. Then $i_x < i_y < j_x$, and therefore, v_{i_y} is adjacent to either v_{i_x} or v_{j_x} . Since v_{i_y} is not adjacent to v_{j_x} , v_{i_y} is adjacent to v_{i_x} . In case when $i_y < i_x$, we also have that v_{i_y} is adjacent to v_{i_x} by the same reason. It implies that $\{v_{i_1}, \dots, v_{i_t}\}$ is a clique. By the symmetric argument, we have that $\{v_{j_1}, \dots, v_{j_t}\}$ is a clique. \diamond

By Claim 4, $\{v_{i_1}, \dots, v_{i_t}\}$ and $\{v_{j_1}, \dots, v_{j_t}\}$ are cliques, and therefore, G contains $K_t \boxminus K_t$ as an induced subgraph, contradiction. We conclude that (T, L) is a linear branch-decomposition of linear mim-width at most $t - 1$. \square

Corollary 3.7. *Let $t \geq 2$ be an integer.*

- (1) *Given an n -vertex $(K_t \boxminus K_t)$ -free co-comparability graph and $\sigma, \rho \subseteq \mathbb{N}$, one can solve MIN-(OR MAX-) (σ, ρ) -DOMINATION in time $\mathcal{O}(n^{3d(t-1)+4})$ where $d = \max(d(\sigma), d(\rho))$.*
- (2) *Given an n -vertex $(K_t \boxminus K_t)$ -free co-comparability graph and a $(q \times q)$ -matrix D_q , one can solve D_q -PARTITIONING in time $\mathcal{O}(qn^{3dq(t-1)+4})$ where $d = \max_{i,j} d(D_q[i, j])$.*




Figure 4: The (4×4) column-clique grid.

Note that permutation graphs do not contain $K_3 \boxplus K_3$ as induced subgraphs, because $K_3 \boxplus K_3$ is the complement of C_6 , which is not a permutation graph. Therefore, Corollary 3.7 generalizes the algorithms for permutation graphs.

In the next, we show that co-comparability graphs have unbounded mim-width. For positive integers p and q , the $(p \times q)$ *column-clique grid* is the graph on the vertex set $\{v_{i,j} : 1 \leq i \leq p, 1 \leq j \leq q\}$ where

- for every $j \in \{1, \dots, q\}$, $\{v_{1,j}, \dots, v_{p,j}\}$ is a clique,
- for every $i \in \{1, \dots, p\}$ and $j_1, j_2 \in \{1, \dots, q\}$, v_{i,j_1} is adjacent to v_{i,j_2} if and only if $|j_2 - j_1| = 1$,
- for $i_1, i_2 \in \{1, \dots, p\}$ and $j_1, j_2 \in \{1, \dots, q\}$ with $i_1 \neq i_2$ and $j_1 \neq j_2$, v_{i_1,j_1} is not adjacent to v_{i_2,j_2} .

We depict an example in Figure 4. For each $1 \leq i \leq p$, we call $\{v_{i,1}, \dots, v_{i,h}\}$ the *i-th row* of G , and define its columns similarly.

Lemma 3.8. *For integers $p, q \geq 12$, the $(p \times q)$ column-clique grid has mim-width at least $\min(\frac{p}{4}, \frac{q}{3})$.*

Proof. Let G be the $(p \times q)$ column-clique grid, and let (T, L) be a branch-decomposition of G . It is enough to show that (T, L) has mim-width at least $\min(\frac{p}{4}, \frac{q}{3})$. Let w be the mim-width of (T, L) . By Lemma 2.3, G admits a vertex bipartition (A, B) where $\text{mim}_G(A) \leq w$ and for each $U \in \{A, B\}$, $\frac{|V(G)|}{3} < |U| \leq \frac{2|V(G)|}{3}$. It is sufficient to show that $\text{mim}_G(A) \geq \min(\frac{p}{4}, \frac{q}{3})$.

Firstly, assume that for each row R of G , $R \cap A \neq \emptyset$ and $R \cap B \neq \emptyset$. Then there is an edge between $R \cap A$ and $R \cap B$, as $G[R]$ is connected. For each i -th row R_i , we choose a pair of vertices $v_{i,a_i} \in R \cap A$ and $v_{i,b_i} \in R \cap B$ that are adjacent. We choose a subset $X \subseteq \{1, \dots, p\}$ such that $|X| \geq \frac{p}{2}$ and every pair (v_{i,a_i}, v_{i,b_i}) in $\{(v_{i,a_i}, v_{i,b_i}) : i \in X\}$ satisfies that $a_i + 1 = b_i$. By taking the same parity of a_i 's, we choose a subset $Y \subseteq X$ such that $|Y| \geq \frac{p}{4}$ and all integers in $\{a_i : i \in Y\}$ have the same parity. Then we can observe that $\{v_{i,a_i}, v_{i,b_i} : i \in Y\}$ is an induced matching in $G[A, B]$, as all integers in $\{a_i : i \in Y\}$ have the same parity. Therefore, we have $\text{mim}_G(A) \geq \frac{p}{4}$.

Now, we assume that there exists a row R such that R is fully contained in one of A and B . Without loss of generality, we may assume that R is contained in A . Since $|B| > \frac{|V(G)|}{3}$, there is a subset $X \subseteq \{1, \dots, q\}$ such that $|X| > \frac{q}{3}$ and for each $i \in X$, the i -th column contains a vertex of B . For each i -th column where $i \in X$, we choose a vertex $v_{a_i,i}$ in B . It is not hard to verify that the edges between $\{v_{a_i,i} : i \in X\}$ and the rows in R form an induced matching of size $\frac{q}{3}$ in $G[A, B]$.

Therefore, we have $d \geq \min(\frac{p}{4}, \frac{q}{3})$. \square

Corollary 3.9. *For every large enough n , there is a co-comparability graph on n vertices having mim-width at least $\sqrt{\frac{n}{12}}$.*

Proof. Let $p \geq 4$ be an integer, and let $n := 12p^2$. Let G be the $(4p \times 3p)$ column-clique grid. It is not hard to see that

$$v_{1,1}, v_{2,1}, \dots, v_{4p,1}, v_{1,2}, v_{2,2}, \dots, v_{4p-1,3p}, v_{4p,3p}$$

is a co-comparability ordering, as each column is a clique. Thus, G is a co-comparability graph. By Lemma 3.8, $\text{mimw}(G) \geq p = \sqrt{\frac{n}{12}}$. \square

We remark that the two classes, $(K_t \boxminus S_t)$ -free chordal graphs and $(K_t \boxminus K_t)$ -free co-comparability graphs, are subclasses of the class of graphs of sim-width at most 1 and having no induced subgraph isomorphic to $K_t \boxminus K_t$ and $K_t \boxminus S_t$ for $t \geq 3$. This is because chordal graphs have no $K_2 \boxminus K_2$ induced subgraph, and co-comparability graphs have no $K_3 \boxminus S_3$ induced subgraphs by Lemma 2.1. Motivated from it, we will extend these classes to classes graphs of bounded sim-width and having no $K_t \boxminus K_t$ and $K_t \boxminus S_t$ as induced subgraphs or induced minors, in Section 4.

4 Graphs of bounded sim-width

In Section 3, we proved that graphs of sim-width at most 1 contain all chordal graphs and all co-comparability graphs. A classical result on chordal graphs is that MINIMUM DOMINATING SET is NP-complete on chordal graphs [6]. So, even for this kind of locally-checkable problems, we cannot expect efficient algorithms on graphs of sim-width at most w . Therefore, to obtain a meta-algorithm for graphs of bounded sim-width encompassing many locally-checkable problems, we must impose some restrictions. We approach this problem in a way analogous to what has previously been done in the realm of rank-width [13].

It is well known that complete graphs have rank-width at most 1, but they have unbounded tree-width. Fomin, Oum, and Thilikos [13] showed that if a graph G is K_r -minor free, then its tree-width is bounded by $c \cdot \text{rw}(G)$ where c is a constant depending on r . This can be utilized algorithmically, to get a result for graphs of bounded rank-width when excluding a fixed minor, as the class of problems solvable in FPT time is strictly larger when parameterized by tree-width than rank-width [20].

We will do something similar by focusing on the distinction between mim-width and sim-width. However, K_r -minor free graphs are too strong, as one can show that on K_r -minor free graphs, the tree-width of a graph is also bounded by some constant factor of its sim-width. To see this, one can use Lemma 4.5 and the result on contraction obstructions for graphs of bounded tree-width [12].

Instead of using minors, we exclude $K_t \boxminus K_t$ and $K_t \boxminus S_t$ as induced subgraphs or induced minors. The induced minor operation is rather natural because sim-width does not increase when taking induced minors; we prove this property in Subsection 4.2. In Subsection 4.3, we prove that chordal graphs having no induced minor isomorphic to $K_3 \boxminus S_3$ have unbounded rank-width. As chordal graphs have no induced minor isomorphic to $K_3 \boxminus K_3$ and they have sim-width 1, this implies that graphs that have bounded sim-width and have no induced minor isomorphic to $K_t \boxminus K_t$ or $K_t \boxminus S_t$ for fixed t have unbounded rank-width. Therefore, such classes still extend classes of bounded rank-width.

We denote by $R(k, \ell)$ the *Ramsey number*, that is the minimum integer satisfying that every graph with at least $R(k, \ell)$ vertices contains either a clique of size k or an independent set of size ℓ . By Ramsey's Theorem [26], $R(k, \ell)$ exists for every pair of positive integers k and ℓ .

4.1 Bounding mim-width

We show the following.

Proposition 4.1. *Every graph with sim-width w and no induced minor isomorphic to $K_t \boxplus K_t$ and $K_t \boxplus S_t$ has mim-width at most $8(w+1)t^3 - 1$.*

Proposition 4.2. *Every graph with sim-width w and no induced subgraph isomorphic to $K_t \boxplus K_t$ and $K_t \boxplus S_t$ has mim-width at most $R(R(w+1, t), R(t, t))$.*

We remark that sometimes this Ramsey number can go down to a polynomial function depending on the underlying graphs. See discussions in Belmonte et al [2].

We first prove Proposition 4.1. We use the following result. Notice that the optimal bound of Theorem 4.3 has been slightly improved by Fox [14], and then by Balogh and Kostochka [1].

Theorem 4.3 (Duchet and Meyniel [10]). *For positive integers k and n , every n -vertex graph contains either an independent set of size k or a K_t -minor where $t \geq \frac{n}{2k-1}$.*

Proof of Proposition 4.1. Let G be a graph with sim-width w and no induced minor isomorphic to $K_t \boxplus K_t$ and $K_t \boxplus S_t$. Let (T, L) be a branch-decomposition of G of sim-width w . Let e be an edge of T and (A, B) be the vertex bipartition of G associated with e . We claim that $\text{mim}_G(A) \leq 8(w+1)t^3 - 1$.

Suppose for contradiction that there is an induced matching $\{v_1w_1, \dots, v_mw_m\}$ in $G[A, B]$ where $v_1, \dots, v_m \in A$, $w_1, \dots, w_m \in B$, and $m \geq 8(w+1)t^3$. Let f be the bijection from $\{v_1, \dots, v_m\}$ to $\{w_1, \dots, w_m\}$ such that $f(v_i) = w_i$ for each $i \in \{1, \dots, m\}$. As $m \geq 8(w+1)t^3$, by Theorem 4.3, the subgraph $G[\{v_1, \dots, v_m\}]$ contains either an independent set of size $2(w+1)t$, or a K_{2t^2} -minor.

First assume that $G[\{v_1, \dots, v_m\}]$ contains a K_{2t^2} -minor. Then there is a minor model of K_{2t^2} in $G[\{v_1, \dots, v_m\}]$, that is, there exist pairwise disjoint subsets S_1, \dots, S_{2t^2} of $\{v_1, \dots, v_m\}$ such that

- for each $i \in \{1, \dots, 2t^2\}$, $G[S_i]$ is connected, and
- for two distinct integers $i, j \in \{1, \dots, 2t^2\}$, there is an edge between S_i and S_j .

For each $i \in \{1, \dots, 2t^2\}$, we choose a representative d_i in each $f(S_i)$ and contract each S_i to a vertex c_i . Let G_1 be the resulting graph. Then $G_1[\{c_1, \dots, c_{2t^2}\}, \{d_1, \dots, d_{2t^2}\}]$ is an induced matching of size $2t^2$, and $\{c_1, \dots, c_{2t^2}\}$ is a clique in G_1 . By the same procedure on $\{d_1, \dots, d_{2t^2}\}$, the subgraph $G_1[\{d_1, \dots, d_{2t^2}\}]$ contains either an independent set of size t , or a K_t -minor. In both cases, one can observe that G_1 contains an induced minor isomorphic to $K_t \boxplus K_t$ or $K_t \boxplus S_t$, hence we obtain a contradiction.

Now assume that $G[\{v_1, \dots, v_m\}]$ contains an independent set $\{c_1, \dots, c_{2(w+1)t}\}$, and for each $i \in \{1, \dots, 2(w+1)t\}$, let $d_i := f(c_i)$. By Theorem 4.3, $G[\{d_1, \dots, d_{2(w+1)t}\}]$ contains either an independent set of size $w+1$ or a K_t -minor. In the former case, we obtain an induced matching of size $w+1$ between A and B in G , contradicting to the assumption that $\text{sim}_G(A) \leq w$. In the latter case, we obtain an induced minor isomorphic to $K_t \boxplus S_t$, contradiction.

We conclude that $\text{mim}_G(A) \leq 8(w+1)t^3 - 1$. Since e is arbitrary, G has mim-width at most $8(w+1)t^3 - 1$. \square

In a similar manner we can prove Proposition 4.2.

Proof of Proposition 4.2. One can easily modify from the proof of Proposition 4.1 by replacing the application of Theorem 4.3 with the Ramsey's Theorem to find a clique or an independent set. \square

As a corollary, we obtain the following. In general, we do not have a generic algorithm to find a decomposition, and thus we assume that the decomposition is given as an input.

Corollary 4.4. *Let $w \geq 1$ and $t \geq 2$ be integers.*

- (1) *Given an n -vertex graph of sim-width at most w and having induced minor isomorphic to neither $K_t \boxplus K_t$ nor $K_t \boxplus S_t$, and $\sigma, \rho \subseteq \mathbb{N}$, one can solve MIN-(OR MAX-) (σ, ρ) -DOMINATION in time $\mathcal{O}(n^{3dt'+4})$ where $d = \max(d(\sigma), d(\rho))$ and $t' = 8(w+1)t^3 - 1$.*
- (2) *Given an n -vertex graph of sim-width at most w and having induced minor isomorphic to neither $K_t \boxplus K_t$ nor $K_t \boxplus S_t$, and a $(q \times q)$ -matrix D_q , one can solve D_q -PARTITIONING in time $\mathcal{O}(qn^{3dq't'+4})$ where $d = \max_{i,j} d(D_q[i, j])$ and $t' = 8(w+1)t^3 - 1$.*
- (3) *Given an n -vertex graph of sim-width at most w and having induced subgraph isomorphic to neither $K_t \boxplus K_t$ nor $K_t \boxplus S_t$, and $\sigma, \rho \subseteq \mathbb{N}$, one can solve MIN-(OR MAX-) (σ, ρ) -DOMINATION in time $\mathcal{O}(n^{3dt'+4})$ where $d = \max(d(\sigma), d(\rho))$ and $t' = R(R(w+1, t), R(t, t))$.*
- (4) *Given an n -vertex graph of sim-width at most w and having induced subgraph isomorphic to neither $K_t \boxplus K_t$ nor $K_t \boxplus S_t$, and a $(q \times q)$ -matrix D_q , one can solve D_q -PARTITIONING in time $\mathcal{O}(qn^{3dq't'+4})$ where $d = \max_{i,j} d(D_q[i, j])$ and $t' = R(R(w+1, t), R(t, t))$.*

4.2 Induced minors

We show that the sim-width of a graph does not increase when taking an induced minor. This is one of the main motivations to consider this parameter.

Lemma 4.5. *The sim-width of a graph does not increase when taking an induced minor.*

Proof. Clearly, the sim-width of a graph does not increase when removing a vertex. We prove for contractions.

Let G be a graph, $v_1v_2 \in E(G)$, and let (T, L) be a branch-decomposition of G of sim-width w for some positive integer w . Let z be the contracted vertex in G/v_1v_2 . We claim that G/v_1v_2 admits a branch-decomposition of G of sim-width at most w . We may assume that G is connected and has at least 3 vertices. For G/v_1v_2 , we obtain a branch-decomposition (T', L') as follows:

- Let T' be the tree obtained from T by removing $L(v_2)$, and smoothing its neighbor.
- Let L' be the function from $V(G/v_1v_2)$ to the set of leaves of T' such that $L'(v) = L(v)$ for $v \in V(G/v_1v_2) \setminus \{z\}$ and $L'(z) = L(v_1)$.

Let e_1 and e_2 be the two edges of T incident with the neighbor of $L(v_2)$, but not incident with $L(v_2)$. Let e_{cont} be the edge of T' obtained by smoothing. By construction, all edges of $E(T') \setminus \{e_{cont}\}$ are contained in T .

Claim 5. *For each $e \in E(T') \setminus \{e_{cont}\}$, the sim_{G/v_1v_2} -width of e in (T', L') is at most the sim_G -width of e in (T, L) .*

Proof. Let $e \in E(T') \setminus \{e_{cont}\}$. Let (A, B) be the vertex bipartition of G/v_1v_2 associated with e . Without loss of generality, we may assume $z \in A$. Suppose there exists an induced matching $\{a_1b_1, \dots, a_mb_m\}$ in G/v_1v_2 with $a_1, \dots, a_m \in A$ and $b_1, \dots, b_m \in B$. Let (A', B') be the vertex bipartition of G associated with e where $v_1 \in A'$. We will show that there is also an induced matching in G of same size between A' and B' .

We have either $v_2 \in A'$ or $v_2 \in B'$. If $z \notin \{a_1, \dots, a_m\}$, then $\{a_1b_1, \dots, a_mb_m\}$ is also an induced matching between A' and B' in G . Without loss of generality, we may assume that $z = a_1$.

Case 1. $v_2 \in A'$.

Proof. In G , one of v_1 and v_2 , say v' , is adjacent to b_1 . Also, v_1 and v_2 are not adjacent to any of $\{a_2, \dots, a_m, b_1, \dots, b_m\}$. Therefore, $\{v'b_1, a_2b_2, \dots, a_mb_m\}$ is an induced matching in G between A' and B' , as required. \square

Case 2. $v_2 \in B'$.

Proof. In this case, $\{v_1v_2, a_2b_2, \dots, a_mb_m\}$ is an induced matching between A' and B' , because v_1 and v_2 are not adjacent to any of $\{a_2, \dots, a_m, b_2, \dots, b_m\}$. \square

It shows that the sim_{G/v_1v_2} -width of e in (T', L') is at most the sim_G -width of e in (T, L) . \diamond

In a similar manner, we can show that the sim_{G/v_1v_2} -width of e_{cont} in (T', L') is at most the minimum of the sim_{G/v_1v_2} -width of e_1 and e_2 in (T, L) . Thus, we conclude that $\text{simw}(G/v_1v_2) \leq \text{simw}(G)$. \square

4.3 Unbounded rank-width

We show that the Hsu-clique chain graph depicted in Figure 5 is chordal, but does not contain $K_2 \boxtimes K_2$ or $K_3 \boxtimes S_3$ as an induced minor. Belmonte and Vatshelle [3, Lemma 16] showed that a $(p \times q)$ Hsu-clique chain graph has rank-width at least $\frac{p}{3}$ when $q = 3p + 1$. So, our algorithmic applications based on Proposition 4.1 or Proposition 4.2 are beyond algorithmic applications of graphs of bounded rank-width.

We formally define Hsu-clique chain graphs. For positive integers p, q , the $(p \times q)$ *Hsu-clique chain grid* is the graph on the vertex set $\{v_{i,j} : 1 \leq i \leq p, 1 \leq j \leq q\}$ where

- for every $j \in \{1, \dots, q\}$, $\{v_{1,j}, \dots, v_{p,j}\}$ is a clique,
- for every $i_1, i_2 \in \{1, \dots, p\}$ and $j \in \{1, \dots, q-1\}$, $v_{i_1,j}$ is adjacent to $v_{i_2,j+1}$ if and only if $i_1 \leq i_2$,
- for $i_1, i_2 \in \{1, \dots, p\}$ and $j_1, j_2 \in \{1, \dots, q\}$, v_{i_1,j_1} is not adjacent to v_{i_2,j_2} if $|j_1 - j_2| > 1$.

Proposition 4.6. *Let $w \geq 1$ and $t \geq 2$ be integers. The class of graphs of sim-width w and having induced minor isomorphic to neither $K_t \boxtimes S_t$ nor $K_t \boxtimes K_t$ has unbounded rank-width.*




Figure 5: The (4×4) Hsu-clique chain graph.

Proof. For this, we provide that Hsu-clique chain grids are chordal and having no induced minor isomorphic to $K_t \boxtimes S_t$ or $K_t \boxtimes K_t$. Since Hsu-clique chain grids have unbounded rank-width [3], it proves the proposition.

Let G be the $(p \times q)$ Hsu-clique chain graph for some positive integers p and q .

First show that G is chordal. Suppose for contradiction that G contains an induced cycle $C = c_1 c_2 \cdots c_m c_1$ where $m \geq 4$. Since each column of G is a clique, all vertices of C are contained in two consecutive columns. But also, since each column contains at most 2 vertices, we have $m = 4$, and two consecutive vertices of C are contained in one column. Without loss of generality, we assume c_1, c_2 are in the i -th column for some $i \in \{1, \dots, q - 1\}$, and c_3, c_4 are in the $(i + 1)$ -th column. This implies that there is an induced matching of size 2 between two columns if we ignore edges in each column. However, this is not possible from the construction. Therefore, G is chordal.

If G contains an induced minor isomorphic to $K_2 \boxtimes K_2$, which is an induced cycle of length 4, then G contains an induced subgraph isomorphic to an induced cycle of length ℓ for some $\ell \geq 4$. This contradicts to the fact that G is chordal. So, G has no induced minor isomorphic to $K_2 \boxtimes K_2$.

We claim that G has no induced minor isomorphic to $K_3 \boxtimes S_3$. For contradiction, suppose there is an induced minor model η of $K_3 \boxtimes S_3$ in G . Let $H := K_3 \boxtimes S_3$. For each $v \in V(H)$, let $I_v = \{j : v_{i,j} \in V(\eta(v))\}$. Let $I := I_{v_1^1} \cup I_{v_2^1} \cup I_{v_3^1}$, and let ℓ and r be the smallest and greatest integers in I , respectively. Let $x, y \in \{v_1^1, v_2^1, v_3^1\}$ such that

1. $V(\eta(x))$ contains a vertex in the ℓ -th column, but for $z \in \{v_1^1, v_2^1, v_3^1\} \setminus \{x\}$, $V(\eta(z))$ has no vertex whose row index is higher than all vertices in $V(\eta(x))$ in the ℓ -th column,
2. similarly, $V(\eta(y))$ contains a vertex in the r -th column, but for $z \in \{v_1^1, v_2^1, v_3^1\} \setminus \{x\}$, $V(\eta(z))$ has no vertex whose row index is lower than all vertices in $V(\eta(y))$ in the r -th column.

As $\{v_1^1, v_2^1, v_3^1\}$ is a clique, it is easy to observe that $I_x \cup I_y = I$. Let ℓ be the integer such that $v_\ell^1 \in \{v_1^1, v_2^1, v_3^1\} \setminus \{x, y\}$. By the choice of x and y , every vertex in $V(\eta(v_\ell^1))$ has a neighbor in $V(\eta(x)) \cup V(\eta(y))$. Thus, it contradicts to the assumption that G contains H as an induced minor. \square

5 Lower bound of sim-width on circle graphs

In this section, we construct a set of circle graphs with unbounded sim-width and no triangle, using the approach similar to the one by Mengel [24]. As $\text{simw}(G) \leq \text{mimw}(G)$ for every graph G , the example also implies that the class of circle graphs has unbounded mim-width.

Theorem 5.1. *Circle graphs have unbounded sim-width, and thus have unbounded mim-width.*




Figure 6: The circle graph G_4 of Lemma 5.4

For an integer $d \geq 0$, a graph G is d -degenerate if every subgraph of G contains a vertex of degree at most d . The following lemma asserts that every d -degenerate graph with large treewidth also has a large mim-width.

Lemma 5.2 (Vatshelle's Thesis [32]; See also Mengel [24]). *For every d -degenerate graph G , $\text{mimw}(G) \geq \frac{\text{tw}(G)+1}{3(d+1)}$.*

Given a bipartite graph with large mim-width, we do not lose much of mim-width after adding edges to one part of the bipartition.

Lemma 5.3 (Mengel [24]). *Let H be a bipartite graph with a bipartition (A, B) , and let G be a graph obtained from H by adding some edges in $H[A]$. Then $\text{mimw}(G) \geq \frac{\text{mimw}(H)}{2}$.*

For an integer $k \geq 1$ and a graph H , a graph G is a k -subdivision of H if G can be built from H by replacing every edge of H with a path of length $k+1$ such that those paths replacing edges of H are internally vertex-disjoint.

Now we prove the main lemma of this section.

Lemma 5.4. *For every integer $k \geq 2$, there is a circle graph G_k satisfying the following.*

1. G_k contains no K_3 as a subgraph.
2. G_k contains a 1-subdivision of a $(k \times k)$ -grid as a subgraph. In particular, $\text{tw}(G_k) \geq k$.
3. $\text{mimw}(G_k) \geq \frac{k+1}{18}$.

Proof. Let H be a 1-subdivision of a $(k \times k)$ -grid. It is easy to see that H is 2-degenerate and has tree-width at least k , since it contains a $(k \times k)$ -grid as a minor. By Lemma 5.2, $\text{mimw}(H) \geq \frac{k+1}{9}$.

For $i, j \in \{1, \dots, k\}$, let $a_{i,j}$ be the vertex on the i -th row and j -th column of the $(k \times k)$ -grid. For $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, k-1\}$, let $h_{i,j}$ be the vertex of degree 2 adjacent to $a_{i,j}$ and $a_{i,j+1}$ in H . For $i \in \{1, \dots, k-1\}$ and $j \in \{1, \dots, k\}$, let $v_{i,j}$ be the vertex of degree 2 adjacent to $a_{i,j}$ and $a_{i+1,j}$ in H . Let us define $A := \{a_{i,j} : i, j \in \{1, \dots, k\}\}$, $B_h := \{h_{i,j} : i \in \{1, \dots, k\}, j \in \{1, \dots, k-1\}\}$, and $B_v := \{v_{i,j} : i \in \{1, \dots, k-1\}, j \in \{1, \dots, k\}\}$. Then H is a bipartite graph with the bipartition $(A, B_h \cup B_v)$. Let $B = B_h \cup B_v$.

Before we describe in detail, we briefly explain how we will construct a circle representation of H . For two points a and b on a circle, we denote by \overline{ab} the chord whose end points are a and b . Let p_1, \dots, p_{2k^2} be distinct $2k^2$ points on the circle in clockwise order. We regard each vertex in A as a chord joining two points p_{2i} and p_{2i-1} for some $i \in \{1, \dots, k^2\}$. Since every vertex in B has a degree 2 in H , the chord representing a vertex in B will be represented by a chord intersecting two disjoint chords $\overline{p_{2i}p_{2i-1}}$ and $\overline{p_{2j}p_{2j-1}}$ for some $i \neq j$ that represents its neighbors. Since the chords representing vertices in B may cross each other, the resulting circle graph will be a graph obtained from H by adding some edges in B . By Lemma 5.3, this graph has mim-width at least $\frac{\text{mimw}(H)}{2} \geq \frac{k+1}{18}$.

We explain how we can assign chords representing vertices in A not to induce K_3 in B .

Let C be a circle and c_1, \dots, c_{6k^2} be distinct $6k^2$ points on C in clockwise order. For $i, j \in \{1, \dots, k\}$, $a_{i,j}$ is represented as follows:

- if i is odd, then $a_{i,j}$ is represented by a chord intersecting $c_{6(i-1)k+6j-5}$ and $c_{6(i-1)k+6j}$,
- if i is even, then $a_{i,j}$ is represented by a chord intersecting $c_{6ik-6j+1}$ and $c_{6ik-6j+6}$.

Remark that we assign chords representing vertices $a_{1,1}, \dots, a_{1,k}, a_{2,k}, \dots, a_{2,1}, \dots$ in clockwise order. We assign chords for vertices in B_h as follows.

- For an odd integer $i \in \{1, \dots, k\}$ and an integer $j \in \{1, \dots, k-1\}$, a chord connecting $c_{6(i-1)k+6j-1}$ and $c_{6(i-1)k+6j+2}$ represents a vertex $h_{i,j} \in B_h$.
- For an even integer $i \in \{1, \dots, k\}$ and an integer $j \in \{1, \dots, k-1\}$, a chord connecting $c_{6ik-6j+2}$ and $c_{6ik-6j-1}$ represents a vertex $h_{i,j} \in B_h$.

We assign chords for vertices in B_v as follows. Note that we assign chords to vertices in B_v so that for $i \in \{1, \dots, k-2\}$ and $j_1, j_2 \in \{1, \dots, k\}$, chords representing two vertices v_{i,j_1} and v_{i+1,j_2} cross if and only if either i is odd and $j_1 < j_2$ or i is even and $j_1 > j_2$.

- For an odd integer $i \in \{1, \dots, k-1\}$ and an integer $j \in \{1, \dots, k\}$, a chord connecting $c_{6(i-1)k+6j-2}$ and $c_{6ik+6(k-j)+3}$ represents a vertex $v_{i,j} \in B_v$.
- For an even integer $i \in \{1, \dots, k-1\}$ and an integer $j \in \{1, \dots, k\}$, a chord connecting $c_{6ik-6j+4}$ and $c_{6ik+6j-3}$ represents a vertex $v_{i,j} \in B_v$.

Let G_k be the circle graph obtained by this procedure. See Figure 6 which describes G_4 . Since every new edge of G_k joins v_{i,j_1} and v_{i+1,j_2} for some $i \in \{1, \dots, k-2\}$ and $j_1, j_2 \in \{1, \dots, k\}$, it is easy to see that G_k contains no K_3 . Note that G_k is a graph obtained from H by adding some edges in B . By Lemma 5.3, the resulting circle graph has mim-width at least $\frac{\text{mimw}(H)}{2} \geq \frac{k+1}{18}$. \square

Proof of Theorem 5.1. By Proposition 4.2, there is an increasing function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that every graph G with no induced subgraph isomorphic to $K_3 \boxminus K_3$ and $K_3 \boxminus S_3$ has mim-width at most $f(\text{simw}(G))$. Since the circle graph G_k of Lemma 5.4 contains no K_3 as a subgraph, we have $f(\text{simw}(G_k)) \geq \text{mimw}(G_k) \geq \frac{k+1}{18}$. It follows that $\text{simw}(G_k)$ should be large for sufficiently large k . \square

6 Concluding remarks

In this paper, we showed that every fixed LC-VSVP problem can be solved in XP time parameterized by t on $(K_t \boxminus S_t)$ -free chordal graphs and $(K_t \boxminus K_t)$ -free co-comparability graphs. We further give a mim-width bound on the class of graphs that have sim-width at most w and do not contain $K_t \boxminus S_t$ and $K_t \boxminus K_t$ as induced minors or induced subgraphs.

Vatshelle [32] asked whether there is a function f such that given a graph G , one can in XP time parameterized by $\text{mimw}(G)$ compute a branch-decomposition of mim-width at most $f(k)$. This still remains as an open problem. We asked the same question for sim-width.

Question 1. *Does there exist a function f such that, given a graph G , one can in XP time parameterized by the sim-width k of G compute a decomposition of sim-width $f(k)$?*

We observed that one cannot expect XP algorithms for MINIMUM DOMINATING SET parameterized by sim-width. However, we know that one can solve MAXIMUM INDEPENDENT SET or 3-COLORING in polynomial time on both chordal graphs and co-comparability graphs, which are all known classes of constant sim-width. We ask whether those problems are NP-complete on graphs of sim-width 1 or not.

Question 2. *Is Maximum Independent Set NP-complete on graphs of sim-width at most 1? Also, is 3-Coloring NP-complete on graphs of sim-width at most 1?*

It would be interesting to find more classes having constant sim-width, but unbounded mim-width. We propose some possible classes, that are also presented in Figure 2.

Question 3. *Do weakly chordal graphs or AT-free graphs have constant sim-width?*

We showed that MINIMUM DOMINATING SET can be solved in time $n^{\mathcal{O}(t)}$ on $(K_t \boxminus S_t)$ -free chordal graphs, but we could not obtain an FPT algorithm. We ask whether it is W[1]-hard or not.

Question 4. *Is MINIMUM DOMINATING SET on chordal graphs W[1]-hard parameterized by the maximum t such that the given graph has an $K_t \boxminus S_t$ induced subgraph?*

References

- [1] J. Balogh and A. Kostochka. Large minors in graphs with given independence number. *Discrete Mathematics*, 311(20):2203 – 2215, 2011.
- [2] R. Belmonte, P. Heggernes, P. van t Hof, A. Rafiey, and R. Saei. Graph classes and ramsey numbers. *Discrete Applied Mathematics*, 173:16 – 27, 2014.
- [3] R. Belmonte and M. Vatshelle. Graph classes with structured neighborhoods and algorithmic applications. *Theoret. Comput. Sci.*, 511:54–65, 2013.

- [4] J. R. S. Blair and B. Peyton. An introduction to chordal graphs and clique trees. In *Graph theory and sparse matrix computation*, volume 56 of *IMA Vol. Math. Appl.*, pages 1–29. Springer, New York, 1993.
- [5] H. L. Bodlaender. A partial k -arboretum of graphs with bounded treewidth. *Theoretical Comput. Sci.*, 209:1–45, 1998.
- [6] K. S. Booth and J. H. Johnson. Dominating sets in chordal graphs. *SIAM J. Comput.*, 11(1):191–199, 1982.
- [7] S. Bova, F. Capelli, S. Mengel, and F. Slivovsky. On compiling cnfs into structured deterministic dnmfs. In *Theory and Applications of Satisfiability Testing - SAT 2015 - 18th International Conference, Austin, TX, USA, September 24-27, 2015, Proceedings*, pages 199–214, 2015.
- [8] B.-M. Bui-Xuan, J. A. Telle, and M. Vatshelle. Fast dynamic programming for locally checkable vertex subset and vertex partitioning problems. *Theoret. Comput. Sci.*, 511:66–76, 2013.
- [9] M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015.
- [10] P. Duchet and H. Meyniel. On hadwiger’s number and the stability number. In B. Bollobas, editor, *Graph Theory Proceedings of the Conference on Graph Theory, Cambridge*, volume 62 of *North-Holland Mathematics Studies*, pages 71 – 73. North-Holland, 1982.
- [11] B. Dushnik and E. W. Miller. Partially ordered sets. *Amer. J. Math.*, 63:600–610, 1941.
- [12] F. V. Fomin, P. Golovach, and D. M. Thilikos. Contraction obstructions for treewidth. *J. Combin. Theory Ser. B*, 101(5):302–314, 2011.
- [13] F. V. Fomin, S. Oum, and D. M. Thilikos. Rank-width and tree-width of H -minor-free graphs. *European J. Combin.*, 31(7):1617–1628, 2010.
- [14] J. Fox. Complete minors and independence number. *SIAM Journal on Discrete Mathematics*, 24(4):1313–1321, 2010.
- [15] S. Gaspers, C. H. Papadimitriou, S. H. Sæther, and J. A. Telle. On satisfiability problems with a linear structure. *Proceedings IPEC 2016, Lecture Notes in Comput. Sci.*, To appear.
- [16] F. Gavril. The intersection graphs of subtrees in trees are exactly the chordal graphs. *J. Combinatorial Theory Ser. B*, 16:47–56, 1974.
- [17] P. Golovach and J. Kratochví l. Computational complexity of generalized domination: a complete dichotomy for chordal graphs. In *Graph-theoretic concepts in computer science*, volume 4769 of *Lecture Notes in Comput. Sci.*, pages 1–11. Springer, Berlin, 2007.
- [18] P. A. Golovach, P. Heggernes, M. M. Kanté, D. Kratsch, S. H. Sæther, and Y. Villanger. *Output-Polynomial Enumeration on Graphs of Bounded (Local) Linear MIM-Width*, pages 248–258. Springer Berlin Heidelberg, Berlin, Heidelberg, 2015.
- [19] P. A. Golovach, J. Kratochví l, and O. r. Suchý. Parameterized complexity of generalized domination problems. *Discrete Appl. Math.*, 160(6):780–792, 2012.

- [20] P. Hliněný, S. Oum, D. Seese, and G. Gottlob. Width parameters beyond tree-width and their applications. *The Computer Journal*, 51(3):326–362, 2008.
- [21] W. L. Hsu and T. H. Ma. Substitution decomposition on chordal graphs and applications. In *ISA '91. Algorithms (Taipei, 1991)*, volume 557 of *Lecture Notes in Comput. Sci.*, pages 52–60. Springer, Berlin, 1991.
- [22] C. G. Lekkerkerker and J. C. Boland. Representation of a finite graph by a set of intervals on the real line. *Fund. Math.*, 51:45–64, 1962/1963.
- [23] R. M. McConnell and J. P. Spinrad. Modular decomposition and transitive orientation. *Discrete Math.*, 201(1-3):189–241, 1999.
- [24] S. Mengel. Lower bounds on the mim-width of some graph classes. *preprint*, arxiv.org/abs/1608.01542, 2016.
- [25] S. Oum and P. Seymour. Approximating clique-width and branch-width. *J. Combin. Theory Ser. B*, 96(4):514–528, 2006.
- [26] F. P. Ramsey. On a problem of formal logic. *Proc. London Math. Soc.*, s2-30:264–286, 1930.
- [27] S. H. Sæther, J. A. Telle, and M. Vatshelle. Solving #sat and MAXSAT by dynamic programming. *J. Artif. Intell. Res. (JAIR)*, 54:59–82, 2015.
- [28] N. Sauer. On the density of families of sets. *Journal of Combinatorial Theory, Series A*, 13(1):145 – 147, 1972.
- [29] P. D. Seymour and R. Thomas. Call routing and the ratcatcher. *Combinatorica*, 14(2):217–241, 1994.
- [30] S. Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific J. Math.*, 41(1):247–261, 1972.
- [31] J. A. Telle and A. Proskurowski. Algorithms for vertex partitioning problems on partial k -trees. *SIAM J. Discrete Math.*, 10(4):529–550, 1997.
- [32] M. Vatshelle. New width parameters of graphs. *Ph.D. thesis, University of Bergen*, 2012.